

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

REQUEST FOR FILING CONTINUATION/DIVISIONAL
APPLICATION UNDER 37 C.F.R. § 1.53(b)

Box PATENT APPLICATION
Assistant Commissioner for Patents
Washington, D.C. 20231

JC542 U.S. PTO
09/287632
04/07/99

Sir:

This is a request for filing a ☒ continuation ☐ divisional application under 37 C.F.R. § 1.53(b) of pending Application No. 09/056,767 filed April 8, 1998 and 09/127,735 filed on August 3, 1998, for METHODS AND MEANS FOR OBTAINING MODIFIED PHENOTYPES, by the following named inventor(s):

- (a) Full Name Peter Michael Waterhouse
- (b) Full Name Ming-Bo Wang
- (c) Full Name Michael Wayne Graham

☒ The entire disclosure of the prior application from which a copy of the oath or declaration is supplied herewith is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

☐ This application is being filed by less than all the inventors named in the prior application. In accordance with 37 C.F.R. 1.63(d)(2), the Commissioner is requested to delete the name(s) of the following person or persons who are not inventors of the invention being claimed in this application.

- (a) Full Name _____
- (b) Full Name _____
- (c) Full Name _____

1. ☐ Enclosed is a copy of the prior Application No. as originally filed on , including copies of the specification, claims, drawings and the executed oath or declaration as filed.
2. ☐ Enclosed is a revised prior application and a copy of the prior executed oath or declaration as filed. No new matter has been added to the revised application.
3. ☐ statement(s) claiming small entity status ☐ are enclosed ☐ were filed in prior Application No. , filed on .

4. ☒ The filing fee is calculated below ☒ and in accordance with the enclosed preliminary amendment:

CLAIMS					
	NO. OF CLAIMS		EXTRA CLAIMS	RATE	FEE
Basic Application Fee					\$760.00
Total Claims	38	MINUS 20 =	18	x \$18.00 =	\$324.00
Independent Claims	12	MINUS 3 =	9	x \$78.00 =	\$702.00
If multiple dependent claims are presented, add \$260.00					
Total Application Fee					\$1,786.00
If small entity status is claimed, subtract 50% of Total Application Fee					
Add Assignment Recording Fee of \$40.00 if Assignment document is enclosed					
TOTAL APPLICATION FEE DUE					\$1,786.00

5. ☐ Charge \$ _____ to Deposit Account No. 02-4800 for the fee due.
6. ☒ A check in the amount of \$ 1,786.00 is enclosed for the fee due.
7. ☒ The Commissioner is hereby authorized to charge any appropriate fees under 37 C.F.R. §§ 1.16, 1.17 and 1.21 that may be required by this paper, and to credit any overpayment, to Deposit Account No. 02-4800. This paper is submitted in triplicate.
8. ☐ Cancel in this application original claims __ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
9. ☒ Amend the specification by inserting before the first line the sentence: --This application is a ☒ continuation, ☐ divisional, of Application No. 09/127,735, filed August 3, 1998 and of 09/056,767, filed April 8, 1998.--
10. ☐ Transfer the drawings from the pending prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate of this paper is enclosed for filing in the prior application file. (May only be used if signed by person authorized under 37 C.F.R. § 1.138 and before payment of issue fee.)

11. ☒ New drawings are enclosed.
12. ☐ Priority of Application No. _____ filed on _____ in _____ (country) is claimed under 35 U.S.C. § 119.
- ☐ The certified copy of the priority application
- ☐ is enclosed
- ☐ was filed on _____ in prior Application No. _____, filed on _____
- ☐ has not yet been filed.
13. ☐ A preliminary amendment is enclosed.
14. ☒ Also enclosed Information Disclosure Statement.
15. ☒ The power of attorney in the prior application is to R. Danny Huntington, Esq..
- a. ☐ The power appears in the original papers in the prior application.
- b. ☒ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- c. ☒ Recognize as Associate Attorney Malcolm K. McGowan, Ph.D., Reg. No. 39,300.
- d. ☒ Address all future communications to: (May only be completed by applicant, or attorney or agent of record.)

R. Danny Huntington, Esq.
BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404

April 7, 1999
Date

By: Malcolm K. McGowan
Malcolm K. McGowan, Ph.D.
Registration No. 39,300

ADDRESS OF
SIGNATOR:

BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404
(703) 836-6620

- ☐ inventor(s)
☐ assignee of complete interest
☐ attorney or agent of record
☒ filed under 37 C.F.R. § 1.34(a)

METHODS AND MEANS FOR OBTAINING MODIFIED PHENOTYPES

1. Field of the Invention

The invention relates to methods for reducing the phenotypic expression of a nucleic acid sequence of interest in eucaryotic cells, particularly plant cells, by simultaneously providing the cells with chimeric genes encoding sense and anti sense RNA molecules comprising nucleotide sequences respectively homologous and complementary to at least part of the nucleotide sequence of the nucleic acid of interest. The sense and antisense RNA molecules may be provided as one RNA molecule, wherein the sense and antisense RNA may be linked by a spacer nucleotide sequence and are capable of forming a double stranded RNA molecule. In one aspect of the invention, the methods are directed towards reducing viral infection, resulting in extreme virus resistance. In another embodiment the methods are directed towards reducing the phenotypic expression of an endogenous gene in a plant cell. The invention further relates to high throughput screening methods for identifying the phenotype endowed by the nucleic acid of interest in plant cells. Also provided are plant cells comprising such RNA molecules, as well as plants consisting essentially of such plant cells.

2. Background of the Invention

In 1985, Sanford and Johnston proposed the concept of parasite-derived resistance. They postulated that key gene products from a parasite expressed in the host in a dysfunctional form, in excess or at a wrong developmental stage, should disrupt the function of the parasite with minimal effect on the host (Sanford & Johnston, 1985). Using the QB bacteriophage as a model, they proposed that expression, in bacteria, of the bacteriophage coat protein or modified replicase or an antisense RNA complementary to the QB genome could all give resistance. They also proposed that such approaches would be applicable, in plants, to plant viruses and particularly the use of a modified plant virus replicase. The expression of the coat protein of the plant virus, tobacco mosaic virus (TMV), in tobacco was the first practical validation of this concept for plant virus resistance. This work (Powell-Abel *et al.*, 1986) showed that the expression of the TMV coat protein, from a transgene under the control of the cauliflower mosaic virus 35S promoter, conferred on the

plants resistance to TMV. The same group (Powell *et al.*, 1990) showed that, generally, plants expressing higher levels of coat protein were more resistant to TMV than plants expressing low levels. Since this demonstration there have been very many examples of plants transformed with virus coat protein genes showing resistance (Table 1). There have also been a number of reports of plant virus resistance in plants expressing wild-type replicase (Braun and Hemenway, 1992, Brederode *et al.*, 1995), truncated replicase (Carr *et al.* 1992), modified replicase (Longstaff *et al.* 1993), or antisense viral RNA (Kawchuck *et al.* 1991).

In 1992, Dougherty and colleagues were using different forms of the coat protein gene of tobacco etch virus (TEV) and discovered that some plants containing untranslatable "sense" coat protein genes and antisense coat protein genes showed extreme resistance (ER) to the virus (Lindbo & Dougherty, 1992 a,b). This resistance was functional at the whole plant level and at the single cell level. TEV was unable to replicate in protoplasts derived from ER plants but replicated to high levels in protoplasts from non-transgenic tobacco. Dougherty *et al.* concluded that the mechanism creating the extreme resistance for the untranslatable sense construct was not the same as the often reported coat protein-mediated strategy. They proposed that the mRNA of the untranslatable sense construct was hybridizing with the minus sense genome of the virus and interfering with the procession of replication complexes on the minus strand. They suggested that the use of viral sequence that could form intramolecular pairing should be avoided as this would interfere with their ability to hybridize to the target viral RNA.

Table 1: Plant species that have been genetically engineered for virus resistance (from Rebecca Grumet, *Hort Science* 30[3] 1995)

Species	Viruses
Tobacco (<i>Nicotiana tabacum</i> L.)	AIMV, ArMV, CMV, PVX, PVY, TEV, TGMV, TMV, TRV, TSV, TSWV
Other Nicotiana spp. (<i>N. debneyii</i> , <i>N. benthamiana</i> , <i>N. clevelandii</i>)	ACMV, BYMV, CyMV, CyRSV, BCTV, PEBV, PPV, PVS, WMV
Potato (<i>Solanum tuberosum</i> L.)	PI, RV, PVY
Tomato (<i>Lycopersicon esculentum</i> L.)	AIMV, CMV, TMV, TYLCV
Cucumber (<i>Cucumis sativus</i> L.)	CMV
Melon (<i>Cucumis melo</i> L.)	CMV, ZYMV
Alfalfa (<i>Medicago sativa</i> L.)	AIMV
Papaya (<i>Carica papaya</i> L.)	PRSV
Corn (<i>Zea mays</i> L.)	MDMV
Rice (<i>Oryza sativa</i> L.)	RSV
Rapeseed (<i>Brassica napus</i> L.)	TYMV

The Dougherty group expanded their investigations to plants containing untranslatable sense potato virus Y (PVY) coat protein genes. They obtained results similar to those with TEV and showed that the plants with ER had high transgene copy number, highly active transcription of the transgenes, and low levels of steady state mRNA from the PVY transgene (Lindbo *et al.* 1993, Smith *et al.* 1994). The following model for this mechanism of the resistance was proposed: the high level of transcription of the viral transgene triggers a cytoplasmic based, post transcriptional cellular surveillance system that targets specific RNAs for elimination. As the transgene encodes a transcript comprising viral sequences the mechanism not only degrades the transgene mRNA but also the same sequences in the viral genomic RNA. A key point in this model is the need for a high level of transcription of the

transgene provided by high copy number (3-8; Goodwin *et al.* 1996). Alternatively, the RNA threshold required to trigger the mechanism can be reached by a more modest transcription level aided by the viral RNA from replication in early infection. This gives rise to a "recovery phenotype" where the plant is initially infected and shows symptoms but then produces new growth without virus symptoms and which are extremely resistant to infection.

This proposal was substantiated by the findings that gene silencing of non-viral transgenes could also be due to a post transcriptional mechanism (Ingelbrecht *et al.* 1994; de Carvalho Niebel *et al.* 1995) operating at an RNA level.

A link between non-viral gene silencing and this pathogen derived resistance was provided by inoculating transgenic plants, in which a GUS transgene was known to be silenced by a post transcriptional mechanism, with a virus containing GUS sequences (English *et al.* 1996). In this situation the plants were extremely resistant to the viral infection. However, the same plants were susceptible to the virus if they contained no GUS sequences.

The degree of viral resistance is not always directly related to the level of viral transgene transcription (Mueller *et al.* 1995; English *et al.* 1996) suggesting that there may be an alternative mechanism of inducing the resistance. To accommodate these discrepancies, an alternative model has been proposed in which the crucial factor affecting the resistance is not the level but the quality of the transgene mRNA (English *et al.* 1996). According to this model, the transgene can only induce resistance (or gene silencing) if it is transcribed to produce "aberrant" RNA. There have been many examples of post-transcriptional gene silencing and methylation of the transgene (Hobbs *et al.* 1990; Ingelbrecht *et al.* 1994) and methylation of the transgene has also been found to be associated in some cases of extreme viral resistance (Smith *et al.* 1994, English 1996). In the proposed model, methylation of the transgene leads to the production of aberrant RNAs which induce the cytoplasmic RNA surveillance system. Baulcombe and English have suggested that this method of induction may be the same as that found for the silencing of *met2* in *A.immersus*. In this system transcription of the *met2* RNA was terminated in the methylated regions of the endogenous gene thus producing aberrant truncated

5 RNAs. It was suggested that the methylation was a consequence of ectopic interaction between the transgene and a homologous region of a corresponding region of the endogenous gene (Barry *et al.* 1993). The presence of multiple transgenes would create an increased likelihood of ectopic pairing and is therefore consistent with the correlation between high copy number and extreme viral resistance (Mueller *et al.*, 1995; Goodwin *et al.* 1996; Pang *et al.*, 1996).

10 This whole area has been reviewed recently (e.g. Baulcombe (1996) and Stam *et al.* (1997)) and several models were presented. All models call for a high degree of sequence specificity because the resistance is very (strain) specific and therefore invoke base pairing interaction with an RNA produced from the transgene. One explanation for the induction of the virus resistance or gene silencing with sense transgenes is that the plant's RNA dependent RNA polymerase generates complementary RNAs using the transgene mRNA as a template (Schiebel *et al.* 1993a,b). This hypothetical complementary RNA (cRNA) has not been detected (Baulcombe 1996) but it is expected that the cRNAs will be small and heterodisperse RNAs rather than full complements (Schiebel 1993ab, Baulcombe 1996) and therefore difficult to detect.

The possible methods of action of the cRNA in mediating the virus resistance or gene silencing (as proposed by Baulcombe, 1996) are :

- 20 1: The cRNA hybridizes with transgene mRNA or viral RNA and the duplex becomes a target for dsRNases.;
- 2: The cRNA hybridizes with target RNA to form a duplex which can arrest translation and consequently have an indirect effect on stability (Green, 1993);
- 25 3: The duplex formed between the cRNA and viral RNA causes hybrid arrest of translation of co-factors required for viral replication; and
4. The hybridization of the cRNA affects intra-molecular base pairing required for viral replication.

30 The current models for virus resistance or gene silencing thus involve the induction of a cytoplasmic surveillance system by either high levels of transgene transcription or by the production of aberrant single stranded mRNA. Once the system is triggered, RNA dependent RNA polymerase makes cRNA from the

transgene mRNA. These cRNAs hybridize to the target RNA either directly affecting its translatability or stability, or marking the RNA for degradation.

US 5,190,131 and EP 0 467 349 A1 describe methods and means to regulate or inhibit gene expression in a cell by incorporating into or associating with the genetic material of the cell a non-native nucleic acid sequence which is transcribed to produce an mRNA which is complementary to and capable of binding to the mRNA produced by the genetic material of that cell.

EP 0 223 399 A1 describes methods to effect useful somatic changes in plants by causing the transcription in the plant cells of negative RNA strands which are substantially complementary to a target RNA strand. The target RNA strand can be a mRNA transcript created in gene expression, a viral RNA, or other RNA present in the plant cells. The negative RNA strand is complementary to at least a portion of the target RNA strand to inhibit its activity *in vivo*.

EP 0 240 208 describes a method to regulate expression of genes encoded for in plant cell genomes, achieved by integration of a gene under the transcriptional control of a promoter which is functional in the host and in which the transcribed strand of DNA is complementary to the strand of DNA that is transcribed from the endogenous gene(s) one wishes to regulate.

EP 0 647 715 A1 and US patents 5, 034,323, 5,231,020 and 5,283,184 describe methods and means for producing plants exhibiting desired phenotypic traits, by selecting transgenotes that comprise a DNA segment operably linked to a promoter, wherein transcription products of the segment are substantially homologous to corresponding transcripts of endogenous genes, particularly endogenous flavonoid biosynthetic pathway genes.

W0 93/23551 describes methods and means for the inhibition of two or more target genes, which comprise introducing into the plant a single control gene which has distinct DNA regions homologous to each of the target genes and a promoter operative in plants adapted to transcribe from such distinct regions RNA that inhibits expression of each of the target genes.

W092/13070 describes a method for the regulation of nucleic acid translation, featuring a responsive RNA molecule which encodes a polypeptide and further

includes a regulatory domain, a substrate region and a ribosome recognition sequence. This responsive RNA molecule has an inhibitor region in the regulatory domain, which regulatory domain is complementary to both a substrate region of the responsive RNA molecule and to an anti-inhibitor region of a signal nucleic acid such that, in the absence of the signal nucleic acid, the inhibitor and substrate regions form a base-paired domain the formation of which reduced the level of translation of one of the protein-coding regions in the responsive RNA molecule compared to the level of translation of that one protein-coding region observed in the presence of the signal nucleic acid.

Metzlaff *et al.*, 1997 describe a model for the RNA-mediated RNA degradation and chalcone synthase A silencing in Petunia, involving cycles of RNA-RNA pairing between complementary sequences followed by endonucleolytic RNA cleavages to describe how RNA degradation is likely to be promoted. Fire *et al.*, 1998 describe specific genetic interference by experimental introduction of double-stranded RNA in *Caenorhabditis elegans*. The importance of these findings for functional genomics is discussed (Wagner and Sun, 1998).

Que *et al.*, 1998 describe distinct patterns of pigment suppression which are produced by allelic sense and antisense chalcone synthase transgenes in petunia flowers and have also analyzed flower color patterns in plants heterozygous for sense and antisense chalcone synthase alleles.

WO 98/05770 discloses antisense RNA with special secondary structures which may be used to inhibit gene expression.

WO 94/18337 discloses transformed plants which have increased or decreased linolenic acids as well as plants which express a linoleic acid desaturase.

US 5,850,026 discloses an endogenous oil from Brassica seeds that contains, after crushing and extracting, greater than 86% oleic acid and less than 2.5% α -linolenic acid. The oil also contains less than 7% linoleic acid. The Brassica seeds are produced by plants that contain seed-specific inhibition of microsomal oleate desaturase and microsomal linoleate desaturase gene expression, wherein the inhibition can be created by cosuppression or antisense technology.

US 5,638,637 discloses vegetable oil from rapeseeds and rapeseed producing the same, the vegetable oil having an unusually high oleic acid content of 80% to 90% by weight based on total fatty acid content.

5

SUMMARY OF THE INVENTION

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211

nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with 10 nt stretch of the complement of the sense nucleotide sequence; wherein the RNA is capable of forming a double stranded RNA region by base-pairing between the regions with sense and antisense nucleotide sequence such that at least the 10 consecutive nucleotides of the sense sequence base pair with the 10 consecutive nucleotides of the antisense sequence, resulting in a(n artificial) hairpin RNA structure.

The invention further provides a method for reducing the gene expression of a gene of interest in plant cells, comprising the step of introducing a first and second chimeric DNA, linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of the transgenic plant cells; wherein the first chimeric DNA comprises a plant-expressible promoter, a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the gene of interest and optionally a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The second chimeric DNA comprises a plant-expressible promoter, a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence and optionally a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The sense and antisense RNA molecules are capable of forming a double stranded, duplex RNA by base-pairing between the regions which are complementary.

Also provided by the invention is a method for obtaining virus resistant organisms, particularly plants, comprising the steps of providing cells of the

organism with a first and second chimeric DNA, wherein the first chimeric DNA comprises a promoter, a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the genome of a virus, capable of infecting the plant and a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The second chimeric DNA comprises a promoter, a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of at least 10 consecutive nucleotides of the sense nucleotide sequence and a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The sense and antisense RNA molecules are capable of forming a double stranded RNA region by base-pairing between the regions which are complementary.

The first and second chimeric DNA are either integrated separately in the nuclear genome of the transformed plant cell or they are linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of the transgenic plant cells.

The invention also provides a method for identifying a phenotype associated with the expression of a nucleic acid of interest in a eucaryotic cell, comprising the steps of selecting within the nucleotide sequence of interest, a target sequence of at least 10 consecutive nucleotides; designing a sense nucleotide sequence corresponding to the length of the selected target sequence and which has a sequence identity of at least about 75% to about 100% with the selected target sequence, designing an antisense nucleotide sequence which has a sequence identity of at least about 75% to about 100% with the complement of the sense nucleotide sequence and comprises a stretch of at least about 10 consecutive nucleotides with 100% sequence identity to the complement of a part of the sense nucleotide sequence. The method further comprises the steps of introducing an RNA molecule comprising both the designed sense and antisense nucleotide

sequences into a suitable eucaryotic host cell comprising the nucleic acid including the nucleotide sequence with hitherto unidentified phenotype; and observing the phenotype by a suitable method.

The invention also provides a eucaryotic cell, comprising a nucleic acid of interest which is normally capable of being phenotypically expressed, further comprising a chimeric DNA molecule comprising a promoter, capable of being expressed in that eucaryotic cell, a DNA region, which when transcribed, yields an RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest and an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence wherein the RNA molecule is capable of forming a double stranded RNA region by base-pairing between the regions with sense and antisense nucleotide sequence such that at least said 10 consecutive nucleotides of the sense sequence base pair with said 10 consecutive nucleotides of the antisense sequence, resulting in a hairpin RNA structure, preferably an artificial hairpin structure and a DNA region involved in transcription termination and polyadenylation, wherein the phenotypic expression of the nucleic acid of interest is significantly reduced.

Also provided by the invention is a eucaryotic cell, comprising a nucleic acid of interest, which is normally capable of being phenotypically expressed, further comprising a chimeric RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest and an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence wherein the RNA is capable of forming a double stranded RNA region by base-pairing between the regions with sense and antisense nucleotide sequence such that at least said 10 consecutive nucleotides of the sense

sequence basepair with said 10 consecutive nucleotides of the antisense sequence, resulting in an artificial hairpin RNA structure.

It is another objective of the invention to provide a eucaryotic cell, comprising a gene of interest, which is normally capable of being phenotypically expressed, further comprising a first and second chimeric DNA, linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of that eucaryotic cell wherein the first chimeric DNA comprises the following operably linked parts a promoter capable of being expressed in the eucaryotic cell a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the gene of interest; and a DNA region involved in transcription termination and polyadenylation; and wherein the second chimeric DNA comprises the following operably linked parts: a promoter operative in the eucaryotic cell; a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence; a DNA region involved in transcription termination and polyadenylation ; wherein the sense and antisense RNA molecules are capable of forming a double stranded RNA region by base-pairing between the regions which are complementary.

It is yet another objective of the invention to provide a virus resistant plant, comprising a first and second chimeric DNA integrated in the nuclear genome of its cells , wherein the first chimeric DNA comprises a plant-expressible promoter, a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the genome of a virus, capable of infecting the plant, and a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The second chimeric DNA comprises a

plant-expressible promoter, a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence, and a DNA region involved in transcription termination and polyadenylation functioning in plant cells. The sense and antisense RNA molecules are capable of forming a double stranded RNA region by base-pairing between the regions which are complementary. The first and second chimeric DNA are integrated either in one locus or in different loci in the nuclear genome.

The invention also provides a method for modifying the fatty acid profile in oil, preferably increasing the oleic acid content, from a plant, preferably from oilseed rape, the method comprising the step of introducing a chimeric DNA into the cells of the plant, comprising the following operably linked parts: a). a plant-expressible promoter, preferably a seed-specific promoter; b). a DNA region, particularly with the nucleotide sequence of SEQ ID No 6, which when transcribed yields an RNA molecule comprising an RNA region capable of forming an artificial stem-loop structure, wherein one of the annealing RNA sequences of the stem-loop structure comprises a nucleotide sequence essentially similar to at least part of the nucleotide sequence of a $\Delta 12$ desaturase encoding open reading frame, and wherein the second of the annealing RNA sequences comprises a sequence essentially similar to at least part of the complement of at least part of the nucleotide sequence of the $\Delta 12$ desaturase encoding open reading frame; and optionally; c) a DNA region involved in transcription termination and polyadenylation. Plants with modified fatty acid profile, particularly with increased oleic acid content, comprising the mentioned chimeric genes are also provided by the invention. Also encompassed are oils obtained from such plants or seed.

With the foregoing and other objects, advantages and features of the invention that will become hereinafter apparent, the nature of the invention may be more clearly understood by reference to the following detailed description of the preferred embodiments of the invention and to the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 represents schematically the different sense and antisense constructs, as well as the so-called CoP (complementary pair) constructs used for obtaining virus resistance (Fig 1B) or for reducing the phenotypic expression of a transgenic Gus gene (Fig 1A).

Figure 2A represents schematically the so-called panhandle construct or CoP constructs used for reducing the phenotypic expression of a $\Delta 12$ desaturase gene in *Arabidopsis* (Nos Pro: nopaline synthase gene promoter; nptII neomycin phosphotransferase coding region; Nos term: nopaline syntase gene terminator; FP1: truncated seed specific napin promoter; 480 bp: 5' end of the Fad2 gene of *Arabidopsis thaliana* in sense orientation; 623 bp: spacer; 480 bp: 5' end of the Fad2 gene of *Arabidopsis thaliana* in antisense orientation).

Figure 2B represents schematically a common cosuppression construct for for reducing the phenotypic expression of a $\Delta 12$ desaturase gene in *Arabidopsis thaliana*.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE INVENTION

One of the objectives of the invention is to provide a eucaryotic cell, particularly a plant cell with an RNA molecule which comprises a hairpin structure including a determined sense part and a determined antisense part. Potentially, an RNA molecule is capable of forming several secondary structures, some of which may contain the desired hairpin. It is expected that the real secondary structure of the RNA in the cell, will have the lowest free energy. In accordance with the invention, the RNA molecule to be produced in the cell is designed in such a way that at least in its lowest free energy state, which it can assume within that cell, it will comprise the desired hairpin.

As used herein "hairpin RNA" refers to any self-annealing double stranded RNA molecule. In its simplest representation, a hairpin RNA consists of a double

stranded stem made up by the annealing RNA strands, connected by a single stranded RNA loop, and is also referred to as a "pan-handle RNA". However, the term "hairpin RNA" is also intended to encompass more complicated secondary RNA structures comprising self-annealing double stranded RNA sequences, but also internal bulges and loops. The specific secondary structure adapted will be determined by the free energy of the RNA molecule, and can be predicted for different situations using appropriate software such as FOLDRNA (Zuker and Stiegler, 1981).

As used herein, "sequence identity" with regard to nucleotide sequences (DNA or RNA), refers to the number of positions with identical nucleotides divided by the number of nucleotides in the shorter of the two sequences. The alignment of the two nucleotide sequences is performed by the Wilbur and Lipmann algorithm (Wilbur and Lipmann, 1983) using a window-size of 20 nucleotides, a word length of 4 nucleotides, and a gap penalty of 4. Computer-assisted analysis and interpretation of sequence data, including sequence alignment as described above, can, e.g., be conveniently performed using the programs of the Intelligenetics™ Suite (Intelligenetics Inc., CA). Sequences are indicated as "essentially similar" when such sequence have a sequence identity of at least about 75%, particularly at least about 80 %, more particularly at least about 85%, quite particularly about 90%, especially about 95%, more especially about 100%, quite especially are identical. It is clear that when RNA sequences are said to be essentially similar or have a certain degree of sequence identity with DNA sequences, thymine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence.

As used herein, the term "plant-expressible promoter" means a DNA sequence which is capable of controlling (initiating) transcription in a plant cell. This includes any promoter of plant origin, but also any promoter of non-plant origin which is capable of directing transcription in a plant cell, *i.e.*, certain promoters of viral or bacterial origin such as the CaMV35S, the subterranean clover virus promoter No 4 or No 7, or T-DNA gene promoters.

The term "expression of a gene" refers to the process wherein a DNA region which is operably linked to appropriate regulatory regions, particularly to a promoter,

is transcribed into an RNA which is biologically active i.e., which is either capable of interaction with another nucleic acid or which is capable of being translated into a polypeptide or protein. A gene is said to encode an RNA when the end product of the expression of the gene is biologically active RNA, such as e.g. an antisense RNA, a ribozyme or a replicative intermediate. A gene is said to encode a protein when the end product of the expression of the gene is a protein or polypeptide.

A nucleic acid of interest is "capable of being expressed", when said nucleic acid, when introduced in a suitable host cell, particularly in a plant cell, can be transcribed (or replicated) to yield an RNA, and/or translated to yield a polypeptide or protein in that host cell.

The term "gene" means any DNA fragment comprising a DNA region (the "transcribed DNA region") that is transcribed into a RNA molecule (e.g., an mRNA) in a cell operably linked to suitable regulatory regions, e.g., a plant-expressible promoter. A gene may thus comprise several operably linked DNA fragments such as a promoter, a 5' leader sequence, a coding region, and a 3' region comprising a polyadenylation site. A plant gene endogenous to a particular plant species (endogenous plant gene) is a gene which is naturally found in that plant species or which can be introduced in that plant species by conventional breeding. A chimeric gene is any gene which is not normally found in a plant species or, alternatively, any gene in which the promoter is not associated in nature with part or all of the transcribed DNA region or with at least one other regulatory region of the gene.

As used herein, "phenotypic expression of a nucleic acid of interest" refers to any quantitative trait associated with the molecular expression of a nucleic acid in a host cell and may thus include the quantity of RNA molecules transcribed or replicated, the quantity of post-transcriptionally modified RNA molecules, the quantity of translated peptides or proteins, the activity of such peptides or proteins.

A "phenotypic trait" associated with the phenotypic expression of a nucleic acid of interest refers to any quantitative or qualitative trait, including the trait mentioned, as well as the direct or indirect effect mediated upon the cell, or the organism containing that cell, by the presence of the RNA molecules, peptide or protein, or posttranslationally modified peptide or protein. The mere presence of a

nucleic acid in a host cell, is not considered a phenotypic expression or a phenotypic trait of that nucleic acid, even though it can be quantitatively or qualitatively traced. Examples of direct or indirect effects mediated on cells or organisms are, e.g., agronomically or industrial useful traits, such as resistance to a pest or disease; higher or modified oil content etc.

As used herein, "reduction of phenotypic expression" refers to the comparison of the phenotypic expression of the nucleic acid of interest to the eucaryotic cell in the presence of the RNA or chimeric genes of the invention, to the phenotypic expression of the nucleic acid of interest in the absence of the RNA or chimeric genes of the invention. The phenotypic expression in the presence of the chimeric RNA of the invention should thus be lower than the phenotypic expression in absence thereof, preferably be only about 25%, particularly only about 10%, more particularly only about 5% of the phenotypic expression in absence of the chimeric RNA, especially the phenotypic expression should be completely inhibited for all practical purposes by the presence of the chimeric RNA or the chimeric gene encoding such an RNA.

A reduction of phenotypic expression of a nucleic acid where the phenotype is a qualitative trait means that in the presence of the chimeric RNA or gene of the invention, the phenotypic trait switches to a different discrete state when compared to a situation in which such RNA or gene is absent. A reduction of phenotypic expression of a nucleic acid may thus, a.o., be measured as a reduction in transcription of (part of) that nucleic acid, a reduction in translation of (part of) that nucleic acid or a reduction in the effect the presence of the transcribed RNA(s) or translated polypeptide(s) have on the eucaryotic cell or the organism, and will ultimately lead to altered phenotypic traits. It is clear that the reduction in phenotypic expression of a nucleic acid of interest, may be accompanied by or correlated to an increase in a phenotypic trait.

As used herein "a nucleic acid of interest" or a "target nucleic acid" refers to any particular RNA molecule or DNA sequence which may be present in a eucaryotic cell, particularly a plant cell.

As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, *e.g.*, a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited ones, *i.e.*, be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA region which is functionally or structurally defined, may comprise additional DNA regions etc.

It has unexpectedly been found by the inventors, that introduction of a chimeric gene capable of being transcribed into an RNA molecule with a nucleotide sequence comprising both the sense and antisense nucleotide sequence of a target gene, or part thereof, integrated in the nuclear genome of a plant cell, could efficiently and specifically reduce the phenotypic expression of that target gene. The reduction in phenotypic expression was more efficient and more predictable than observed when separate chimeric genes were introduced in similar cells with the target gene, encoding either sense or antisense RNA molecules.

At the same time, it has also been found that simultaneously introducing separate chimeric genes in one cell encoding RNA molecules with nucleotide sequences comprising sense and antisense respectively, resulted in extreme virus resistance, even when the chimeric genes were transcribed from weaker promoters. It is well known that gene silencing and virus resistance can be mediated by similar phenomena.

In one embodiment of the invention, a method for reducing the phenotypic expression of a nucleic acid of interest, which is normally capable of being expressed in a eucaryotic cell, particularly a plant cell, is provided, comprising the steps of introducing a chimeric DNA comprising the following operably linked parts:

- a) a promoter, operative in that cell, particularly a plant-expressible promoter;
- b) a DNA region capable of being transcribed into an RNA molecule with a nucleotide sequence comprising

- i. a sense nucleotide sequence of at least 10 nt, preferably 15 nt consecutive nucleotides having between 75 and 100% sequence

identity with at least part of the nucleotide sequence of the nucleic acid of interest; and

- ii. an antisense nucleotide sequence including at least 10 , preferably 15 nt consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 , preferably 15 nt consecutive nucleotides of the sense nucleotide sequence;

wherein the RNA is capable of forming a double stranded RNA by base-pairing between the regions with sense and antisense nucleotide sequence resulting in a hairpin RNA structure; and

- c) a DNA region involved in transcription termination and polyadenylation functioning in the suitable eucaryotic cells, particularly functioning in plant cells.

In a preferred embodiment of the invention, the RNA molecule transcribed from the chimeric gene, consists essentially of the hairpin RNA.

The order of the sense and antisense nucleotide sequence in the RNA molecule is thought not be critical.

Thus, in other words, the chimeric DNA has a transcribed DNA region, which when transcribed, yields an RNA molecule comprising an RNA region capable of forming an artificial stem-loop structure, wherein one of the annealing RNA sequences of the stem-loop structure comprises a sequence, essentially similar to at least part of the nucleotide sequence of the nucleic acid of interest, and wherein the second of the annealing RNA sequences comprises a sequence essentially similar to at least part of the complement of at least part of the nucleotide sequence of the nucleic acid of interest.

The RNA molecule may comprise several artificial hairpin structures, which may be designed to reduce the phenotypic expression of different nucleic acids of interest.

In one preferred embodiment, the nucleic acid of interest, whose phenotypic expression is targeted to be reduced, is a gene incorporated in the genome of a eucaryotic cell, particularly a plant cell. It will be appreciated that the means and methods of the invention can be used for the reduction of phenotypic expression of

a gene which belongs to the genome of the cell as naturally occurring, (an endogenous gene), as well as for the reduction of phenotypic expression of a gene which does not belong to the genome of the cell as naturally occurring, but has been introduced in that cell (a transgene). The transgene can be introduced stably or transiently, and can be integrated into the nuclear genome of the cell, or be present on a replicating vector, such as a viral vector.

In another preferred embodiment, the nucleic acid of interest, whose phenotypic expression is targeted to be reduced is a viral nucleic acid, particularly a viral RNA molecule, capable of infecting a eucaryotic cell, particularly a plant cell. In this case, the phenotype to be reduced is the replication of the virus, and ultimately, the disease symptoms caused by the infecting virus.

Preferably, the nucleotide sequence of the target nucleic acid corresponding to the sense nucleotide sequence is part of a DNA region which is transcribed, particularly a DNA region which is transcribed and translated (in other words a coding region). It is particularly preferred that the target sequence corresponds to one or more consecutive exons, more particularly is located within a single exon of a coding region.

The length of the sense nucleotide sequence may vary from about 10 nucleotides (nt) up to a length equaling the length (in nucleotides) of the target nucleic acid. Preferably the total length of the sense nucleotide sequence is at least 10 nt, preferably 15 nt, particularly at least about 50 nt, more particularly at least about 100 nt, especially at least about 150 nt, more especially at least about 200 nt, quite especially at least about 550 nt. It is expected that there is no upper limit to the total length of the sense nucleotide sequence, other than the total length of the target nucleic acid. However for practical reason (such as e.g. stability of the chimeric genes) it is expected that the length of the sense nucleotide sequence should not exceed 5000 nt, particularly should not exceed 2500 nt and could be limited to about 1000 nt.

It will be appreciated that the longer the total length of the sense nucleotide sequence is, the less stringent the requirements for sequence identity between the total sense nucleotide sequence and the corresponding sequence in the target gene

become. Preferably, the total sense nucleotide sequence should have a sequence identity of at least about 75% with the corresponding target sequence, particularly at least about 80 %, more particularly at least about 85%, quite particularly about 90%, especially about 95%, more especially about 100%, quite especially be identical to the corresponding part of the target nucleic acid. However, it is preferred that the sense nucleotide sequence always includes a sequence of about 10 consecutive nucleotides, particularly about 20 nt, more particularly about 50 nt, especially about 100 nt, quite especially about 150 nt with 100% sequence identity to the corresponding part of the target nucleic acid. Preferably, for calculating the sequence identity and designing the corresponding sense sequence, the number of gaps should be minimized, particularly for the shorter sense sequences.

The length of the antisense nucleotide sequence is largely determined by the length of the sense nucleotide sequence, and will preferably correspond to the length of the latter sequence. However, it is possible to use an antisense sequence which differs in length by about 10%. Similarly, the nucleotide sequence of the antisense region is largely determined by the nucleotide sequence of the sense region, and preferably is identical to the complement of the nucleotide sequence of the sense region. Particularly with longer antisense regions, it is however possible to use antisense sequences with lower sequence identity to the complement of the sense nucleotide sequence, preferably with at least about 75% sequence identity, more preferably with at least about 80%, particularly with at least about 85%, more particularly with at least about 90% sequence identity, especially with at least about 95% sequence to the complement of the sense nucleotide sequence. Nevertheless, it is preferred that the antisense nucleotide sequences always includes a sequence of about 10 , preferably 15 consecutive nucleotides, particularly about 20 nt, more particularly about 50 nt, especially about 100 nt, quite especially about 150 nt with 100% sequence identity to the complement of a corresponding part of the sense nucleotide sequence. It is clear that the length of the stretch of the consecutive nucleotides with 100% sequence identity to the complement of the sense nucleotide sequence cannot be longer than the sense nucleotide sequence itself. Again, preferably the number of gaps should be minimized, particularly for the shorter

antisense sequences. Further, it is also preferred that the antisense sequence has between about 75% to 100% sequence identity with the complement of the target sequence.

The RNA molecule resulting from the transcription of the chimeric DNA may comprise a spacer nucleotide sequence located between the sense and antisense nucleotide sequence. In the absence of such a spacer sequence, the RNA molecule will still be able to form a double-stranded RNA, particularly if the sense and antisense nucleotide sequence are larger than about 10 nucleotides and part of the sense and/or antisense nucleotide sequence will be used to form the loop allowing the base-pairing between the regions with sense and antisense nucleotide sequence and formation of a double stranded RNA. It is expected that there are no length limits or sequence requirements associated with the spacer region, as long as these parameters do not interfere with the capability of the RNA regions with the sense and antisense nucleotide sequence to form a double stranded RNA. In a preferred embodiment, the spacer region varies in length from 4 to about 200 bp, but as previously mentioned, it may be absent.

In a preferred embodiment, the hairpin RNA formed by the sense and antisense region and if appropriate the spacer region, is an artificial hairpin RNA. By "artificial hairpin RNA" or "artificial stem-loop RNA structure", is meant that such hairpin RNA is not naturally occurring in nature, because the sense and antisense regions as defined are not naturally occurring simultaneously in one RNA molecule, or the sense and antisense regions are separated by a spacer region which is heterologous with respect to the target gene, particularly, the nucleotide sequence of the spacer has a sequence identity of less than 75% with the nucleotide sequence of the target sequence, at the corresponding location 5' or 3' of the endpoints of the sense nucleotide sequence. A hairpin RNA can also be indicated as artificial, if it is not comprised within the RNA molecule it is normally associated with. It is conceivable to use in accordance with the invention a chimeric DNA whose transcription results in a hairpin RNA structure with a naturally occurring nucleotide sequence (which otherwise meets the limits as set forth in this specification)

provided this hairpin RNA is devoid of the surrounding RNA sequences (not involved in the hairpin structure formation).

Although it is preferred that the RNA molecule comprising the hairpin RNA does not further comprise an intron sequence, it is clear that the chimeric DNA genes encoding such RNAs may comprise in their transcribed region one or more introns.

In fact, the inventors have unexpectedly found that inclusion of an intron sequence in the chimeric DNA genes encoding an RNA molecule comprising the hairpin RNA, preferably in the spacer region, and preferably in sense orientation, enhances the efficiency of reduction of expression of the target nucleic acid. The enhancement in efficiency may be expressed as an increase in the frequency of plants wherein silencing occurs or as an increase in the level of reduction of the phenotypic trait. In a particularly preferred embodiment, the intron is essentially identical in sequence to the *Flaveria trinervia* pyruvate orthophosphate dikinase 2 intron 2, more particularly, it comprises the sequence of SEQ ID No 7.

It has been observed that contrary to methods using either antisense or sense nucleotide sequences alone to reduce the phenotypic expression of a target nucleic acid (which generally depend on the dosage of sense or antisense molecule, and thus the chimeric genes encoding the sense and antisense molecules need to be under the control of relatively strong promoters) the method of the current invention does not rely on the presence of such strong promoter regions to drive the transcriptional production of the RNA comprising both the sense and antisense region. In other words, a whole range of promoters, particularly plant expressible promoters, is available to direct the transcription of the chimeric genes of the invention. These include, but are not limited to strong promoters such as CaMV35S promoters (e.g., Harpster et al., 1988). In the light of the existence of variant forms of the CaMV35S promoter, as known by the skilled artisan, the object of the invention can equally be achieved by employing these alternative CaMV35S promoters and variants. It is also clear that other plant-expressible promoters, particularly constitutive promoters, such as the opine synthase promoters of the *Agrobacterium* Ti- or Ri-plasmids, particularly a nopaline synthase promoter, or

subterranean clover virus promoters can be used to obtain similar effects. Also contemplated by the invention are chimeric genes to reduce the phenotypic expression of a nucleic acid in a cell, which are under the control of single subunit bacteriophage RNA polymerase specific promoters, such as a T7 or a T3 specific promoter, provided that the host cells also comprise the corresponding RNA polymerase in an active form.

It is a further object of the invention, to provide methods for reducing the phenotypic expression of a nucleic acid in specific cells, particularly specific plant cells by placing the chimeric genes of the invention under control of tissue-specific or organ-specific promoters. Such tissue-specific or organ-specific promoters are well known in the art and include but are not limited to seed-specific promoters (e.g., WO89/03887), organ-primordia specific promoters (An *et al.*, 1996), stem-specific promoters (Keller *et al.*, 1988), leaf specific promoters (Hudspeth *et al.*, 1989), mesophyll-specific promoters (such as the light-inducible Rubisco promoters), root-specific promoters (Keller *et al.*, 1989), tuber-specific promoters (Keil *et al.*, 1989), vascular tissue specific promoters (Peleman *et al.*, 1989), stamen-selective promoters (WO 89/10396, WO 92/13956), dehiscence zone specific promoters (WO 97/13865) and the like.

In another embodiment of the invention, the expression of a chimeric gene to reduce the phenotypic expression of a target nucleic acid can be controlled at will by the application of an appropriate chemical inducer, by operably linking the transcribed DNA region of the chimeric genes of the invention to a promoter whose expression is induced by a chemical compound, such as the promoter of the gene disclosed in European Patent publication ("EP") 0332104, or the promoter of the gene disclosed in WO 90/08826.

It has been found that a similar reduction in phenotypic expression of a nucleic acid of interest in a eucaryotic cell, particularly in a plant cell, can be obtained by providing the sense and antisense RNA encoding transcribable DNA regions as separate transgenes, preferably located in one locus, particularly as one allele.

Thus, in another embodiment of the invention a method for reducing the phenotypic expression of a nucleic acid which is normally capable of being expressed in a eucaryotic cell, particularly a plant cell, is provided, comprising the steps of introducing a first and second chimeric DNA, linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of the transgenic cells;

wherein the first chimeric DNA comprises the following operably linked parts:

- a) a promoter, operative in the cell, particularly a plant-expressible promoter;
- b) a DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10, preferably 15 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and
- c) a DNA region involved in transcription termination and polyadenylation functioning in the corresponding eucaryotic cell; and

wherein the second chimeric DNA comprises the following operably linked parts:

- a) a promoter, operative in the cell, particularly a plant-expressible promoter;
- b) a DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10, preferably 15 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10, preferably 15 consecutive nucleotides of the sense nucleotide sequence;
- c) a DNA region involved in transcription termination and polyadenylation functioning in the corresponding eucaryotic cell;

wherein the sense and antisense RNA are capable of forming a double stranded RNA by base-pairing between the regions which are complementary.

Preferred embodiments for the different structural and functional characteristics, such as length and sequence of sense, antisense and spacer regions, of this method are as described elsewhere in this specification.

The RNA molecule, comprising the sense and antisense nucleotide sequences capable of forming a hairpin structure, which are produced by the transcription of the chimeric genes, can also be introduced directly in a plant cell to reduce the phenotypic expression of the target nucleic acid, particularly to reduce the phenotypic expression of a targeted endogenous gene, or a targeted transgene. Such RNA molecules could be produced e.g. by

1. cloning the DNA region capable of being transcribed into an RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest and an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence, whereby the RNA is capable of forming a double stranded RNA by base-pairing between the regions with sense and antisense nucleotide sequence resulting in a hairpin RNA structure, under control of a promoter suitable for recognition by a DNA-dependent RNA polymerase in an *in vitro* transcription reaction, such as but not limited to a T7-polymerase specific promoter;
2. performing an *in vitro* transcription reaction by adding *inter alia* the suitable DNA -dependent RNA polymerase as well as the required reagents to generate the RNA molecules; and
3. isolating the RNA molecules.

In vitro transcription methods as well as other methods for *in vitro* RNA production are well known in the art and commercial kits are available. Methods for direct introduction of RNA in plant cells are also available to the skilled person and include but are not limited to electroporation, microinjection and the like.

The chimeric gene(s) for reduction of the phenotypic expression of a target nucleic acid of interest in a cell, may be introduced either transiently, or may be stably integrated in the nuclear genome of the cell. In one embodiment, the chimeric

gene is located on a viral vector, which is capable of replicating in the eucaryotic cell, particularly the plant cell (see e.g., WO 95/34668 and WO 93/03161).

The recombinant DNA comprising the chimeric gene to reduce the phenotypic expression of a nucleic acid of interest in a host cell, may be accompanied by a chimeric marker gene, particularly when the stable integration of the transgene in the genome of the host cell is envisioned. The chimeric marker gene can comprise a marker DNA that is operably linked at its 5' end to a promoter, functioning in the host cell of interest, particularly a plant-expressible promoter, preferably a constitutive promoter, such as the CaMV 35S promoter, or a light inducible promoter such as the promoter of the gene encoding the small subunit of Rubisco; and operably linked at its 3' end to suitable plant transcription 3' end formation and polyadenylation signals. It is expected that the choice of the marker DNA is not critical, and any suitable marker DNA can be used. For example, a marker DNA can encode a protein that provides a distinguishable color to the transformed plant cell, such as the A1 gene (Meyer *et al.*, 1987), can provide herbicide resistance to the transformed plant cell, such as the *bar* gene, encoding resistance to phosphinothricin (EP 0,242,246), or can provide antibiotic resistance to the transformed cells, such as the *aac(6')* gene, encoding resistance to gentamycin (WO94/01560).

A recombinant DNA comprising a chimeric gene to reduce the phenotypic expression of a gene of interest, can be stably incorporated in the nuclear genome of a cell of a plant. Gene transfer can be carried out with a vector that is a disarmed Ti-plasmid, comprising a chimeric gene of the invention, and carried by *Agrobacterium*. This transformation can be carried out using the procedures described, for example, in EP 0 116 718.

Alternatively, any type of vector can be used to transform the plant cell, applying methods such as direct gene transfer (as described, for example, in EP 0 233 247), pollen-mediated transformation (as described, for example, in EP 0 270 356, WO85/01856 and US 4,684,611), plant RNA virus-mediated transformation (as described, for example, in EP 0 067 553 and US 4,407,956), liposome-mediated transformation (as described, for example, in US 4,536,475), and the like.

Other methods, such as microprojectile bombardment as described for corn by Fromm *et al.* (1990) and Gordon-Kamm *et al.* (1990), are suitable as well. Cells of monocotyledonous plants, such as the major cereals, can also be transformed using wounded and/or enzyme-degraded compact embryogenic tissue capable of forming compact embryogenic callus, or wounded and/or degraded immature embryos as described in WO92/09696. The resulting transformed plant cell can then be used to regenerate a transformed plant in a conventional manner.

The obtained transformed plant can be used in a conventional breeding scheme to produce more transformed plants with the same characteristics or to introduce the chimeric gene for reduction of the phenotypic expression of a nucleic acid of interest of the invention in other varieties of the same or related plant species, or in hybrid plants. Seeds obtained from the transformed plants contain the chimeric genes of the invention as a stable genomic insert.

It is a further object of the invention to provide eucaryotic cells, preferably plant cells and organisms (preferably plants) comprising the chimeric genes for the reduction of the phenotypic expression of a target nucleic acid as described in the invention.

It is a yet a further object of the invention to provide plant cells, comprising a nucleic acid of interest, which is normally capable of being expressed phenotypically, further comprising an RNA molecule with a nucleotide sequence which includes:

- i. a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and
- ii. an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of the at least 10 consecutive nucleotides of the sense nucleotide sequence and capable of forming a double stranded RNA by association with the sense nucleotide sequence;

wherein the phenotypic expression of the nucleotide acid of interest is significantly reduced by the presence of the RNA molecule, when compared to the phenotypic expression of the nucleic acid of interest in the absence of the RNA molecule.

The RNA molecule may be encoded by chimeric DNA. Preferred embodiments for the sense and antisense nucleotide sequence, particularly concerning length and sequence, are as mentioned elsewhere in this specification.

It will be appreciated that the methods and means described in the specification can also be applied in High Throughput Screening (HTS) methods, for the identification or confirmation of phenotypes associated with the expression of a nucleic acid sequence with hitherto unidentified function in a eucaryotic cell, particularly in a plant cell.

Such a method comprises the steps of

1. selecting a target sequence within the nucleic acid sequence of interest with unidentified or non-confirmed function/phenotype when expressed. Preferably, if the nucleic acid has putative open reading frames, the target sequence should comprise at least part of one of these open reading frames. The length of the target nucleotide sequence may vary from about 10 nucleotides up to a length equaling the length (in nucleotides) of the nucleic acid of interest with unidentified function.
2. designing an RNA molecule comprising sense nucleotide sequence and antisense nucleotide sequence in accordance with the invention.
3. introducing the RNA molecule comprising both the sense and antisense nucleotide sequences designed on the basis of the target sequence, into a suited host cell, particularly a plant cell, comprising the nucleic acid with the nucleotide sequence with hitherto unidentified phenotype. The RNA can either be introduced directly, or can be introduced by means of a chimeric DNA comprising a promoter operative in the host cell of interest, particularly a plant-expressible promoter, and a DNA region functioning as a suitable 3' end formation and polyadenylation signal (terminator) functioning in the host cell, with in-between a DNA region which can be transcribed to yield the RNA molecule comprising the sense and antisense nucleotide sequence. The chimeric DNA can either be introduced transiently or

integrated in the nuclear genome . The chimeric DNA can also be provided on a viral vector (see, e.g., WO 95/34668 and WO 93/03161)

4. observing the phenotype by a suitable method. Depending on the phenotype expected, it may be sufficient to observe or measure the phenotype in a single cell, but it may also be required to culture the cells to obtain an (organized) multicellular level, or even to regenerate a transgenic organism, particularly a transgenic plant.

In its most straightforward embodiment, the RNA molecule comprising both the sense and antisense nucleotide sequences to at least part of a nucleic acid of interest, suitable for the methods of the invention, can be obtained by cloning two copies of a DNA region with the selected target sequence in inverted repeat orientation (preferably separated by a short DNA region which does not contain a transcription termination signal, and encodes the spacer sequence) under a suitable promoter. This chimeric DNA is then either used as template DNA in an *in vitro* transcription method to generate the RNA molecule, which is introduced in the host cell, or the chimeric DNA itself is introduced in the host cell.

The methods and means of the invention can thus be used to reduce phenotypic expression of a nucleic acid in a eucaryotic cell or organism, particularly a plant cell or plant, for obtaining shatter resistance (WO 97/13865), for obtaining modified flower color patterns (EP 522 880, US 5,231,020), for obtaining nematode resistant plants (WO 92/21757, WO 93/10251, WO 94/17194), for delaying fruit ripening (WO 91/16440, WO 91/05865, WO 91/16426, WO 92/17596, WO 93/07275, WO 92/04456, US 5,545,366), for obtaining male sterility (WO 94/29465, WO89/10396, WO 92/18625), for reducing the presence of unwanted (secondary) metabolites in organisms, such as glucosinolates (WO97/16559) or chlorophyll content (EP 779 364) in plants , for modifying the profile of metabolites synthesized in a eucaryotic cell or organisms by metabolic engineering e.g. by reducing the expression of particular genes involved in carbohydrate metabolism (WO 92/11375, WO 92/11376, US 5, 365, 016, WO 95/07355) or lipid biosynthesis (WO 94/18337, US 5, 530, 192), for delaying senescence (WO 95/07993), for altering lignification in plants (WO 93/05159, WO 93/05160), for altering the fibre quality in cotton (US 5,

597, 718), for increasing bruising resistance in potatoes by reducing polyphenoloxidase (WO 94/03607), etc.

The methods of the invention will lead to better results and/or higher efficiencies when compared to the methods using conventional sense or antisense nucleotide sequences and it is believed that other sequence-specific mechanisms regulating the phenotypic expression of target nucleic acids might be involved and/or triggered by the presence of the double-stranded RNA molecules described in this specification.

A particular application for reduction of the phenotypic expression of a transgene in a plant cell, *inter alia*, by antisense or sense methods, has been described for the restoration of male fertility, the latter being obtained by introduction of a transgene comprising a male sterility DNA (WO 94/09143, WO 91/02069). The nucleic acid of interest is specifically the male sterility DNA. Again, the processes and products described in this invention can be applied to these methods in order to arrive at a more efficient restoration of male fertility.

The methods and means of the invention, particularly those involving RNA molecules comprising a hairpin RNA and the encoding chimeric genes, have proven to be particularly suited for the modification of the composition of oil content in plants, particularly in seeds. Particularly preferred plants are crop plants used for oil production such as but not limited to oilseed rape (*Brassica juncea*, *napus*, *rapa*, *oleracea*, *campestris*), corn, cotton, groundnut, sunflower, castor beans, flax, coconut, linseed, soybean. Preferred target genes are the desaturase genes, particularly $\Delta 12$ desaturase encoding genes such as those encoded by the Fad2 genes, especially the genes whose nucleotide sequence can be found in the Genbank Database under accession number AF123460 (from *Brassica carinata*), AF12042841 (*Brassica rapa*), L26296 (*Arabidopsis thaliana*) or A65102 (*Corylus avellana*). It is clear that it is well within the reach of the person skilled in the art to obtain genes homologous to the disclosed fad2 genes from other species e.g. by hybridization and/or PCR techniques.

Preferred embodiments for the configuration of sense and antisense nucleotide sequences, particularly concerning length and sequence are as

mentioned elsewhere in this specification. Also, preferred embodiments for chimeric genes encoding hairpin containing RNA molecules, particularly concerning promoter elements are as described elsewhere in the specification. For this application, it is particularly preferred that the promoters are seed-specific promoters.

5 In a preferred embodiment, the artificial hairpin RNA comprising RNA molecule thus comprises part of a $\Delta 12$ desaturase encoding ORF in sense orientation and a similar part in antisense orientation, preferably separated by a spacer sequence. In a particularly preferred embodiment the artificial hairpin RNA (or its encoding chimeric gene) comprises the nucleotide sequence of SEQ ID No 6
10 or a similarly constructed nucleotide sequence based on the aforementioned Brassica fad2 genes.

Preferably the chimeric gene encoding the artificial hairpin RNA is transcribed under control of a seed-specific promoter, particularly under control of the FPI promoter as described elsewhere in this application.

15 A reduction of the expression of $\Delta 12$ desaturase gene in oil containing plants leads to increase in oleic acid and a concomitant decrease in linolenic acid and linoleic acid. A higher frequency of plants with oil wherein the increase in oleic acid and concomitant decrease in linolenic and linoleic acid is significant is found using the means and methods of the invention than in transgenic plants harboring
20 common cosuppression constructs. Moreover the absolute levels of increase, respectively decrease are higher respectively lower than in transgenic plants harboring common cosuppression constructs.

Using the means and methods of the invention, it is thus possible to obtain plants and seeds, comprising an oil of which the composition after crushing and
25 extracting has an increased oleic acid content (expressed as a percentage of the total fatty acid composition), particularly a three fold increase, when compared with control plants.

It is expected that using the methods and means of the invention, transgenic Brassica plant can be obtained, whose seeds comprise an oil wherein the oleic acid
30 content exceeds 90% of the total fatty acid contents.

The methods and means of the invention will be especially suited for obtaining virus resistance, particularly extreme virus resistance, in eucaryotic cells or organisms, particularly in plant cells and plants. A non-limiting list of viruses for plants against which resistance can be obtained, is represented in Table I.

5 The methods and means of the invention further allow the use of viral genes, hitherto unused for obtaining virus resistant plants in addition to the commonly used coat protein genes or replicase genes. Such different viral genes include protease encoded genes (Pro) genome linked protein (Vpg) encoding genes, cytoplasmic inclusion body encoding genes (CI) as target nucleic acid sequences for obtaining
10 virus resistant plants.

It is clear that the invention will be especially suited for the reduction of phenotypic expression of genes belonging to multigene families.

It is also clear that the methods and means of the invention are suited for the reduction of the phenotypic expression of a nucleic acid in all plant cells of all plants,
15 whether they are monocotyledonous or dicotyledonous plants, particularly crop plants such as but not limited to corn, rice, wheat, barley, sugarcane, cotton, oilseed rape, soybean, vegetables (including chicory, brassica vegetables, lettuce, tomato), tobacco, potato, and sugarbeet, but also plants used in horticulture, floriculture or forestry. The means and methods of the invention will be particularly suited for
20 plants which have complex genomes, such as polyploid plants.

It is expected that the chimeric RNA molecules produced by transcription of the chimeric genes described herein, can spread systemically throughout a plant, and thus it is possible to reduce the phenotypic expression of a nucleic acid in cells of a non-transgenic scion of a plant grafted onto a transgenic stock comprising the
25 chimeric genes of the invention (or vice versa) a method which may be important in horticulture, viticulture or in fruit production.

The following non-limiting Examples describe the construction of chimeric genes for the reduction of the phenotypic expression of a nucleic acid of interest in a eucaryotic cell and the use of such genes. Unless stated otherwise in the
30 Examples, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory*

Manual, Second Edition, Cold Spring Harbor Laboratory Press, NY and in Volumes 1 and 2 of Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, *Current Protocols*, USA. Standard materials and methods for plant molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

Throughout the description and Examples, reference is made to the following sequences:

SEQ ID N° 1: sequence of the Potato virus Y fragment of the Nia gene used for the construction of various sense and antisense constructs, for obtaining virus resistance.

SEQ ID N° 2: sequence of the coding region of the Gusd CoP construct of Example 1.

SEQ ID N° 3: sequence of a modified 5' untranslated region (5'UTR) from Johnsongrass mosaic virus

SEQ ID N° 4: sequence of the Subterranean clover virus promoter No 4 with S7 enhancer.

SEQ ID N° 5: sequence of the Subterranean clover virus double enhancer promoter No 4.

SEQ ID N° 6: sequence of the CoP construct for the $\Delta 12$ desaturase gene expression inhibition.

SEQ ID N° 7 : sequence of the *Flaveria trinervia* pyruvate orthophosphate dikinase intron

The following free text is included in the sequence listing :

<223> fragment of the NIa ORF

<223> Description of Artificial Sequence: coding region of the Gusd CoP construct

<223> deficient Gus coding region

<223> antisense to the 5' end of the Gus coding region

<223> Description of Artificial Sequence: 5'UTR of Johnson mosaic virus

<223> Description of Artificial Sequence: Subterranean clover virus S4 promoter with S7 enhancer

<223> Description of Artificial Sequence: subterranean clover virus promoter S4 with S4 enhancer

<223> Description of Artificial Sequence: coding sequence of the desaturase CoP construct

<223> corresponding to the 5' end of the delta12-desaturase (fad2) coding region, in sense orientation

<223> corresponding to the 5' end of the delta12-desaturase (fad2) coding region, in anti sense orientation

<223> Description of Artificial Sequence: intron 2 of the Flaveria trinervia purvate orthophosphate dikinase

The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

EXAMPLE 1

Experimental procedures

Gene construction

Standard gene cloning methods (Sambrook *et al.* 1989) were used to make the chimeric genes. A schematic representation of the constructs used is shown in Figures 1A and 1B.

The components for these constructs were:

- * Cauliflower mosaic virus 35S promoter from the Cabb-JI isolate(35S) (Harpster *et al.*, 1988)
- * Octopine synthase terminator (ocs-t) (MacDonald *et al.*, 1991)
- * Subterranean clover virus promoter No 4 (S4) (WO 9606932)
- * Subterranean clover virus terminator No 4 (s4t) (WO 9606932)
- * Subterranean clover virus double enhancer promoter No 4 (S4S4) (SEQ ID N° 5)
- * Subterranean clover virus promoter No 4 with S7 enhancer (S7S4) (SEQ ID N° 4)
- * Maize ubiquitin promoter (Ubi) (Christensen and Quail, 1996)
- * Agrobacterium tumour morphology 1 gene terminator (tm1') (Zheng *et al.* , 1991)
- * the Nia gene of an Australian strain of Potato virus Y (Nia) (SEQ ID N° 1)

* a dysfunctional β -glucuronidase open reading frame encoding DNA (Gusd)(SEQ ID N° 2 from nucleotide 1 to nucleotide 1581)

* a modified 5' untranslated region (5'UTR) from Johnsongrass mosaic virus (JGMV5') (SEQ ID N° 3).

5 This contains insertion of a *NcoI* site at the ATG start codon followed by three stop codons in frame, and a *PstI* site (for insertion of the intron as in constructs 4 and 5 of Figure 1A). In vector constructs 2 and 6 of Figure 1A, the Gusd open reading frame is inserted in at the *NcoI* site, removing the stop codons; in all other constructs of Fig 1A it is inserted downstream of the *PstI* site.

10 * a castor bean catalase intron (Ohta et al., 1990) as modified by Wang *et al.*(1997) ("intron").

The chimeric genes were constructed by operably assembling the different parts as schematically indicated in Figure 1A or Figure 1B and inserting the obtained chimeric genes in the T-DNA vectors pART27 and pART7 vectors (Gleave, 1992) between the left T-DNA border and the chimeric plant expressible *neo* gene.

15 The DNA encoding a dysfunctional β -glucuronidase open reading frame (GUSd) was obtained by deleting from a *gus* coding region the sequence between the two *EcoRV* restriction sites. For the construction of the chimeric gene encoding the RNA molecule comprising both sense and antisense nucleotide sequence to a β -glucuronidase gene, a sequence was added to the Gusd gene to be allow base pairing the 5'end over 558 bases resulting in a sequence as represented in SEQ ID N° 2. This sequence was cloned between the maize ubiquitin promoter and the tm1' terminator and inserted in a T-DNA vector.

20 T-DNA vectors were constructed which comprised a first and a second chimeric virus resistance gene, wherein the first chimeric gene consisted of:

- 25 1. a CaMV 35S promoter sequence, coupled to
2. in sense orientation, the nucleotide sequence from PVY encoding either
 - Vpg protein (see e.g., Genbank Accession Nr ZZ9526 from nucleotide 1013 to nucleotide 1583), or
 - 30 • part of the CI protein (see e.g., Genbank Accession Nr M95491 from nucleotide 3688 to nucleotide 4215) or

- Protease (Pro) (see e.g., EMBL Accession Nr D00441 from nucleotide 5714 to nucleotide 7009), followed by
- 3. the S4 terminator from subterranean clover mosaic virus, as described above.

5 The second chimeric gene consists of

1. a S4 promoter as described above, coupled to
2. in anti-sense orientation, the nucleotide sequence from PVY encoding either
 - Vpg protein , or
 - 10 • CI protein or
 - Protease , followed by
3. the octopine synthase terminator as described above.

The sense and antisense sequences within one T-DNA vector were derived from the same PVY coding region.

15 Also, T-DNA vectors were constructed for use in altering the fatty acid composition in oil (see Fig 2), comprising

1. a FP1 promoter (truncated seed specific napin promoter, containing sequences between -309 and +1, as described in Stalberg et al; linked to
2. the nucleotide sequence of SEQ ID No 6, comprising the 480 bp located 5' in the ORF encoding the $\Delta 12$ desaturase from *Arabidopsis thaliana* (Fad2) in sense orientation and in antisense orientation, linked by a 623 bp spacer sequence; followed by
- 20 3. the terminator from the nopaline synthase gene.

In addition, T-DNA vectors were constructed to evaluate the influence of a presence of an intron sequence in the chimeric genes encoding CoP constructs. To this end, constructs were made comprising:

1. a CamV35S promoter, followed by
2. the protease encoding ORF from PVY (see above) in sense orientation;
3. the sequence of SEQ ID No 7 (encoding the *Flaveria trinervia* pyruvate orthophosphate dikinase intron 2)
- 30 4. the protease encoding ORF from PVY in antisense orientation; and

5. the octopine synthase gene terminator.

Plant transformation

Nicotiana tabacum (W38) tissue was transformed and regenerated into whole plants essentially as described by Landsman *et al.* (1988). Rice (*Oryza sativa*) was transformed essentially as described by Wang *et al.* (1997).

Rice supertransformation

Mature embryos from a rice plant expressing GUS and hygromycin phosphotransferase (HPT) activity were excised from mature seed and placed on callus inducing media for 7 weeks. Calli were recovered from these cultures, incubated with *Agrobacteria* containing various binary vector constructs for 2 days, then placed on callusing media containing hygromycin, bialaphos and Timentin™. During the next four weeks hygromycin and bialaphos resistant calli developed. These callus lines were maintained on hygromycin and bialaphos containing media for a further 2 months before being assayed for GUS activity.

GUS Assay

Rice calli were tested for GUS activity using the histochemical stain X-glucuronide or the fluorogenic substrate 4-methyl-umbeliferone glucuronide (MUG) essentially as described by Jefferson *et al.* (1987).

EXAMPLE 1. Comparison of chimeric genes comprising only antisense, only sense, or both sense and antisense (Complimentary Pair (CoP)) sequence for reduction in phenotypic expression of an integrated β -glucuronidase gene.

Transgenic rice tissue expressing β -glucuronidase (GUS) from a single transgene (and hygromycin resistance from a *hph* gene) (lines V10-28 and V10-67) was supertransformed using vectors that contained the *bar* gene conferring phosphinothricin resistance and various sense, antisense and CoP constructs (see Figure 1A) derived from a crippled GUS (GUSd) gene. The supertransformed tissue was maintained on hygromycin and bialaphos selection media for 3 weeks then

analyzed for GUS activity. A crippled GUS gene was used so that expression from this gene would not be superimposed on the endogenous GUS activity.

The figures in Table 2 represent the rate of MU production measured by absorption at 455 nm, with excitation at 365 nm of 1.5 µg of total protein in a reaction volume of 200 µl. The rate was measured over 30 min at 37°C. The reading for non-transgenic rice calli was 0.162. The figures in bracket which follow the description of the introduced construct refer to Figure 1A.

The results (Table 2) showed that supertransformation with the binary vector containing the *bar* gene without the GUSd gene had no silencing effect on the endogenous GUS activity. Supertransformation with GUSd in a sense or antisense orientation, with or without an intron or an early stop codon, showed some degree of reduction (in about 25% of the analyzed calli) of the endogenous GUS activity (see last two rows in Table 2 representing the percentage of analyzed calli with a MUG assay reading of less than 2.000). However, supertransformation with a CoP construct gave in about 75% to 100% of the analyzed calli, reduction of the endogenous GUS activity. This CoP construct was designed so that the 3' end of the mRNA produced could form a duplex with the 5' end of the transcript to give a "pan-handle" structure .

These data show that a complimentary pair can be made using one self-annealing transcript, that this design is much more effective than a conventional sense or antisense construct, and that the approach can be used to reduce the phenotypic expression of genes present in a plant cell.

Table 2. MUG assay of Supertransformed Rice Calli

	Vector cassette (1)	Sense (2)	Sense + Stop (3)	Sense + stop+ intron (4)	Antisense + stop + intron (5)	Inverted repeat CoP (6)
V10-28	121,0	97,45	38,43	38,88	<u>0,290</u>	<u>0,565</u>
	45,58	6,637	64,16	115,5	<u>0,572</u>	<u>0,316</u>
	99,28	71,60	149,2	133,0	37,2	<u>0,351</u>
	26,17	<u>0,224</u>	<u>0,955</u>	98,46	53,94	<u>0,210</u>
	92,21	<u>0,321</u>	68,32	<u>0,502</u>	105,5	<u>0,701</u>
	108,8	5,290	105,6	39,35	56,73	<u>0,733</u>
	6,432	<u>0,9460</u>	136,6	<u>1,545</u>	60,36	2,103
	90,80	32,44	140,4	10,36	71,12	119,8
	98,24	128,8	62,38	111,6	13,17	<u>0,717</u>
	93,76	31,28	17,79	14,42	<u>0,424</u>	<u>0,398</u>
		5,023		88,06	26,98	<u>0,315</u>
		40,27		52,28	115,5	<u>0,270</u>
		36,40		30,26	149,7	16,78
		53,24		107,5	66,75	67,28
		29,97		26,75	145,8	<u>0,217</u>
		89,06		105,1	<u>0,534</u>	<u>0,208</u>
		<u>0,256</u>		135,1	9,4	
		68,23		95,04	35,33	
		5,481		71,5		
V10-67	318,8	93,43	<u>0,199</u>	31,82	<u>1,395</u>	<u>0,472</u>
	109,5	73,19	<u>0,197</u>	58,08	152,4	<u>0,256</u>
	30,35	128,1	<u>0,157</u>	56,32	67,42	<u>0,296</u>
	40,04	<u>1,506</u>	128	44,62	12,11	<u>0,452</u>
	228	140,6	130,3	<u>0,454</u>	<u>0,668</u>	<u>0,422</u>
	23,05	<u>1,275</u>	196,2	17,32	23,34	<u>0,196</u>
	241,2	<u>0,272</u>	12,43	73,2	76,10	<u>0,294</u>
	118,5	<u>0,209</u>	140,0	20,32	130,1	<u>0,172</u>
	11,27	42,05	90,13	107,4	<u>0,841</u>	<u>0,436</u>
	110,6	117,5	157,4	<u>0,453</u>	66,12	<u>0,398</u>
	19,29	118,9	<u>0,518</u>	87,81	136,9	<u>0,242</u>
	121,0	21,44	<u>0,231</u>	<u>0,299</u>	67,92	
	115,1	155,0	116,1	<u>0,206</u>	50,32	
	77,1	190,9	43,18	12,47	170,3	
	106,1	<u>0,773</u>	31,06	<u>0,213</u>	108,9	
	73,12	<u>0,146</u>		11,15	<u>1,241</u>	
	29,97			19,22	4,092	
	50,11				169,6	
	80,34				76,88	
	117,8				22,08	
	159,1				91,6	
	67,52				7,855	
	92,32				69,76	
	27,97				<u>0,822</u>	
V10-28	0%	21%	10%	10,5%	22%	75%
V10-67	0%	37,5%	33%	29,5%	21%	100%

EXAMPLE 2. Comparison of the efficiency of using chimeric genes comprising only antisense genes, only sense genes, or both genes simultaneously for obtaining virus resistance in transgenic plants.

5 Gene constructs were made using the PVY protease encoding sequence (SEQ ID N° 1) in a sense orientation, an antisense orientation and in a complimentary pair (CoP) orientation, where the T-DNA comprised both the sense and antisense chimeric genes each under control of their own promoter. In all three arrangements the CaMV35S promoter was used. Five different versions of CoP
10 constructs were made in which the second promoter was either the CaMV35S promoter, the S4 promoter, the double S4 promoter, the S7 enhanced S4 promoter, or the vascular specific *rolC* promoter (see Figure 1B).

15 These constructs were transformed into tobacco (via *Agrobacterium* mediated DNA transfer) and approximately 25 independently transformed plants were recovered per chimeric gene construct. The transgenic plants were transferred to soil and maintained in the greenhouse. About 1 month after transplanting to soil, the plants were inoculated manually with potato virus Y, using standard application methods. Two and four weeks later the plants were scored for virus symptoms. The results (Table 3) showed that after 1 month, 2 on 27 plants comprising only the
20 sense gene, and 1 on 25 plants comprising only the antisense gene showed no symptoms.

25 In contrast respectively 11 on 24 (35S-Nia/S4-antisenseNia construct), 7 on 25 (35S-Nia/RolC-antisenseNia construct), 10 on 27 (35S-Nia/35S-antisenseNia), 7 on 26 (35S-Nia/S4S4-antisenseNia construct), and 7 on 25 (35S-Nia/S7S4-antisenseNia construct) plants which contained both the sense and antisense genes, showed no symptoms. Plants that showed no symptoms were considered to be showing extreme resistance to PVY. They continued to show no symptoms for a further 2 months of monitoring (indicated as Extreme Resistant (ER) in Table 3). Some other plants, particularly those containing CoP constructs, showed a delay
30 and restriction of symptoms. They showed no symptoms 2 weeks after inoculation but showed some minor lesions in some plants after 4 weeks. These plants were

clearly much less effected by PVY than non-transgenic or susceptible tobaccos and were scored as resistant (indicated as ER* in Table 3).

Table 3 Resistance to PVY infection of transgenic tobacco plants comprising either the sense chimeric PVY protease construct, the antisense chimeric PVY protease construct, or both (different CoP constructs).

Sense gene	Antisense gene	Extreme Resistant plants (ER)	"Resistant" plants (ER*)	Total number of transgenic plants
35S-Nia		2	2	27
	35S-AntisenseNia	1	0	25
35S-Nia	35S- AntisenseNia	10	2	27
35S-Nia	S4-AntisenseNia	11	2	24
35S-Nia	RolC-AntisenseNia	7	3	25
35S-Nia	S4S4-AntisenseNia	7	7	26
35S-Nia	S7S4-AntisenseNia	7	4	25

The data show that using CoP constructs results in a much higher frequency of transgenic plants with extreme resistance and resistance than by using either sense or antisense constructs alone.

Next, the copy number of the transgenes in the virus resistant transgenic plants was determined. Therefore, DNA was extracted from all the transgenic plants showing extreme resistance or resistance. DNA was also extracted from five susceptible plants for each construct. The DNA was examined for gene copy number using Southern analysis. The data (Table 4) showed that the genomes of some of the CoP plants showing extreme resistance, particularly the 35S-Nia/S4-AntisenseNia plants, only contained a single copy of the gene construct.

Table 4. Copy number of transgenes comprising sense chimeric PVY protease construct, the antisense chimeric PVY protease construct, or both (different CoP constructs) in extreme resistant, resistant and susceptible plants.

Sense gene	Antisense gene	Extreme Resistant plants (ER)	"Resistant" plants (ER*)	Susceptible plants
35S-Nia		6	1	1/1/1/1/1
	35S-AntisenseNia	4	-	1/8/0/2/1
35S-Nia	35S-AntisenseNia	3/1/2/3/6/3/2/4/2/3	1/1	1/2/2/6/1
35S-Nia	S4-AntisenseNia	2/4/1/3/2/4/6/1/1/3/1	2/6	5/8/2/3
35S-Nia	RolC-AntisenseNia	6/7/6/7/7/7/6	2/1	2/2/2/1/2
35S-Nia	S4S4-AntisenseNia	1/2/4/5/2/2/2	1/1/2/1/1/1/1	1/1/7/1/1
35S-Nia	S7S4-AntisenseNia	2/4/12/5/2/2/7	3/2/1/2	1/1/3/1/1

EXAMPLE 3. Inheritance of extreme resistance in plants from Example 2.

Plants from Example 2 were allowed to self-fertilize and their seeds were collected. Seeds originating from plants showing extreme resistance and low transgene copy number for CoP constructs 35S-Nia/S4-AntisenseNia and 35S-Nia/35S-AntisenseNia, and seeds from the sense and the antisense plants showing extreme resistance, were germinated and grown in the glasshouse. Plants were also grown from seed collected from two susceptible CoP lines, two susceptible sense gene only lines and two susceptible antisense gene only lines. Twenty plants from each line were selected for overall uniformity of size and development stage, put into individual pots, allowed to recover for one week, then inoculated with PVY. The plants were scored for virus symptoms 2,4, and 7 weeks after inoculation. The results (Table 5) showed that all eight plant lines of 35S-Nia/S4-antisenseNia and 35S-Nia/35S-antisenseNia containing one or two gene copies showed an about 3:1 segregation ratio of extreme resistance : susceptible. The progeny of the single antisense gene only line that had given extreme resistance at T_0 , and the progeny of the extremely resistant sense plant containing one gene copy, gave abnormal segregation ratios (2: 18; ER:susceptible). The progeny of the one sense plant that gave extreme resistance and contained 6 gene copies gave a ~ 3:1 ratio (ER:

susceptible). All the progeny of the susceptible T₀ plants showed complete susceptibility to PVY.

These data show extreme resistance from CoP constructs gives stable expression of the resistance which is inherited in a Mendelian way. This also indicates that, in these lines the PVY CoP gene loci are ~100% effective at conferring extreme resistance whereas the transgene loci in the antisense line and one of the two sense lines are only partially effective at conferring extreme resistance.

Table 5.

	35S Sense Nia and S4 Antisense Nia		35S Sense Nia and S4 Antisense Nia		35S Sense Nia		35S Antisense Nia	
	Copy N°	T1 ER:S	Copy N°	T1 ER:S	Copy N°	T1 ER:S	Copy N°	T1 ER:S
ER plant 1	1	15:5	1	16:4	6	17:3	4	2:18
2	1	12:8	2	15:5	1	2:18		
3	1	14:6	2	16:4				
4	1	15:5	2	16:4				
Susceptible Plant 1	8	0:20	1	0:20	1	0:20	1	0:20
2	2	0:20	1	0:20	1	0:20	8	0:20

EXAMPLE 4: Extreme virus resistant transgenic tobacco with different components (sense gene and antisense gene) in different loci within the transgenic plant.

PVY susceptible plants containing the sense transgene (which contained single transgene copies; see Table 6) were crossed with PVY susceptible plants containing the antisense transgene (which had also been analyzed for copy number by Southern analysis; see Table 6). Twenty of the resulting progeny per cross were propagated in the glasshouse, then inoculated with PVY and scored for virus infection as described in Example 2. The progeny from the crosses (between single genes/loci containing plants) would be expected to be in the following ratio: ¼ sense

gene alone, $\frac{1}{4}$ antisense gene alone, $\frac{1}{4}$ comprising both sense and antisense genes, and $\frac{1}{4}$ comprising no genes at all. The results (Table 4) show that, with one exception, a proportion of the progeny from all the successful crosses showed extreme resistance, whereas none of the progeny from selfed sense or selfed antisense plants showed extreme resistance. The one cross that gave no extremely resistant progeny was derived from the parent plant Antisense 2 (As2) which contained 8 copies of the antisense gene. All twenty progeny plants from crosses Sense 1 (S1)(male) \times Antisense 1 (As1) (female) and Sense 3 (S3) (female) \times Antisense 4 (As4) (male) were examined by Southern analysis. The results showed that in both crosses, the plants that showed extreme resistance (or in one case resistance) contained both the sense and antisense genes, whereas plants with no transgenes (nulls), or sense or antisense genes alone, were all susceptible to PVY. To further confirm this absolute correlation between the presence of a complimentary pair (sense with antisense genes) within a plant and extreme resistance, all progeny plants showing extreme resistance were analyzed by Southern blots. The results showed that every extremely resistant or resistant plant contained both sense and antisense genes.

These data show that a complimentary pair gives resistance or extreme resistance even when the genes encoding the sense and antisense genes are not co-located in the genome. The "complimentary pair phenomenon" is not simply due to increased transgene dosage as it would be expected that $\frac{1}{4}$ of the selfed progeny would be homozygous and thus have double the gene dosage, yet they were susceptible.

Table 6: PVY resistance of the progeny plants resulting from crosses between susceptible transgenic tobacco plants comprising the 35S-senseNia gene (S-lines) and susceptible transgenic tobacco plants comprising the 35S-antisenseNia gene (As-lines).

Male parent	Female parent	Extreme Resistant plants (ER)	"Resistant" plants (ER*)
S1	As1	8	
S1	As4	6	5
S2	As4	1	3
S3	As4	3	3
S4	As1	7	1
S3	As5	1	2
S4	As2	0	
S4	As4	2	0
S5	As4	9	4
S5	As5	2	3
As4	As4	0	
As5	As5	0	
S2	S2	0	
S4	S4	0	

Extreme resistant plants showed no symptoms of PVY infection after 7 weeks.

Resistant plants showed very minor lesions 7 weeks after PVY infection.

S1, S2, S3, S4 and S5 are PVY susceptible transgenic tobacco plants comprising the 35S-senseNia gene construct which all have one copy of the transgene integrated.

As1, As2, As4 and As5 are PVY susceptible transgenic tobacco plants comprising the 35S-antisenseNia gene construct which have respectively 1, 8, 2 and 1 copies of the transgene integrated.

EXAMPLE 5. Evaluation of the use of different viral genes as target nucleic acid sequences in obtaining extreme virus resistant genes.

The T-DNA vectors comprising first and second chimeric virus resistance genes based on sequences derived from the coding region for protease, Vpg or CI proteins from PVY as described in this application, were used to obtain transformed tobacco plants, which were subsequently challenged with PVY. The results are summarized in the following table:

Table 7.

Construct	Number of immune plants/ Number of independent transgenic plants	
	Replica 1	Replica 2
35S-Pro sense /S4-Pro antisense	11/24	7/25
35S-Vpg sense /S4-Vpg antisense	8/20	6/18
35S-CI sense /S4-CI antisense	2/23	1/20

Example 6. Intron enhanced silencing

The T-DNA vectors comprising the chimeric genes encoding the CoP constructs wherein an intron (*Flaveria trinervia* pyruvate orthophosphate dikinase intron 2) has been inserted in either the sense orientation or the antisense orientation, between the sense and antisense sequences corresponding to the protease encoding ORF from PVY (as described elsewhere in this application) were used to obtain transformed tobacco plants, which were subsequently challenged with PVY. The results are summarized in the following table:

Table 8.

Construct	Number of immune plants/ Number of independent transgenic plants
35S-Pro(sense)-intron(sense)-Pro(antisense)-Ocs-t	22/24
35S-Pro(sense)-intron(antisense)-Pro(antisense)-Ocs-t	21/24

EXAMPLE 7. Modifying oil profile using CoP constructs in *Arabidopsis*.

T-DNA vectors for modifying the fatty acid composition in oil, extracted from crushed seeds as described elsewhere in this application were used to introduce the chimeric gene encoding the CoP construct for reducing the expression (see Fig 2A; SEQ ID No 6) the $\Delta 12$ desaturase gene (Fad2) in *Arabidopsis thaliana*.

For comparison of the efficiency, transgenic *Arabidopsis* plants were generated wherein the Fad2 gene expression was reduced by a plain cosuppression construct, comprising the FPI seed-specific promoter coupled to the complete ORF from the $\Delta 12$ desaturase gene (Fad2) in *Arabidopsis thaliana* and the nopaline synthase promoter (see Fig 2B).

As control plants, transgenic *Arabidopsis* transformed by unrelated T-DNA constructs were used.

Seeds were harvested, crushed and extracted and the percentage of the major fatty acids in the oil was determined by methods available in the art. The results, which are the mean of two readings, are summarized in Table 9.

Table 9.

Sample	Peak Names ----->												
Name	Myristic	Palmitic	Palmitoleic	Stearic	Oleic	Linoleic	Linolenic	20:0	20:1	22:0	22:1	24:0	C18:1/(C18:2 + C18:3)
Hairpin 1.1	0.00	6.06	0.52	3.21	56.65	7.50	6.82	1.46	16.02	0.00	1.76	0.00	3.95
Hairpin 1.2	0.12	6.86	0.39	3.40	51.28	10.00	8.73	1.64	15.60	0.00	1.97	0.00	2.74
Hairpin 1.3	0.11	8.47	0.50	3.49	21.64	28.99	18.51	2.02	14.19	0.00	2.09	0.00	0.46
Hairpin 1.4	0.00	6.14	0.50	3.37	51.70	9.77	8.02	1.73	16.04	0.00	2.05	0.67	2.91
Hairpin 2.1	0.06	5.19	0.43	3.33	54.84	5.52	7.76	1.77	18.50	0.34	1.83	0.45	4.13
Hairpin 2.2	0.04	7.67	0.46	3.75	19.60	28.29	18.64	2.55	15.96	0.19	2.28	0.56	0.42
Hairpin 3.1	0.00	7.99	0.53	3.62	19.52	28.41	19.24	2.32	15.14	0.00	2.23	0.99	0.41
Hairpin 3.2	0.09	7.00	0.54	3.69	49.02	11.03	9.64	1.71	14.94	0.00	1.72	0.62	2.37
Hairpin 3.3	0.00	5.68	0.49	3.98	46.19	12.82	9.71	2.10	16.70	0.00	1.94	0.39	2.05
Hairpin 3.4	0.17	7.19	0.77	3.69	45.90	11.86	10.65	1.84	15.39	0.00	1.90	0.65	2.04
Hairpin 3.5	0.00	6.45	0.48	3.26	51.76	8.13	10.04	1.51	16.08	0.00	1.92	0.36	2.85
Hairpin 3.6	0.08	7.51	0.23	3.59	19.97	29.13	20.12	2.15	14.54	0.29	2.02	0.36	0.41
Hairpin 3.7	0.14	7.20	0.78	2.90	26.37	24.81	17.18	1.92	15.50	0.36	2.30	0.53	0.63
Hairpin 3.8	0.11	6.34	0.46	3.23	38.58	15.25	13.54	1.89	16.91	0.00	2.36	1.34	1.34
Hairpin 3.9	0.00	6.47	0.49	3.32	47.59	11.44	9.63	1.68	15.96	0.00	1.88	1.55	2.26
Hairpin 3.10	0.00	6.77	0.56	3.48	53.30	7.57	9.34	1.55	15.65	0.00	1.79	0.00	3.15
Hairpin 3.11	0.00	7.05	0.59	3.61	53.62	8.87	8.36	1.55	14.35	0.00	1.99	0.00	3.11

Sample	Peak Names ----->												
Name	Myristic	Palmitic	Palmitoleic	Stearic	Oleic	Linoleic	Linolenic	20:0	20:1	22:0	22:1	24:0	C18:1/(C18:2 + C18:3)
Cosuppresion 1.3	0.07	6.65	0.40	3.42	38.34	15.25	14.16	1.91	17.19	0.31	1.94	0.35	1.30
Cosuppresion 1.4	0.00	8.22	0.57	3.82	18.27	28.82	19.63	2.56	14.83	0.00	2.46	0.83	0.38
Cosuppresion 1.5	0.00	7.51	0.52	3.84	34.59	17.90	14.64	2.18	16.27	0.00	2.02	0.54	1.06
Cosuppresion 1.6	0.07	7.44	0.47	3.16	23.97	27.32	17.29	2.03	15.52	0.18	2.22	0.33	0.54
Cosuppresion 2.1	0.07	7.46	0.43	3.00	23.91	27.21	17.79	1.84	15.27	0.30	2.14	0.58	0.53
Cosuppresion 2.2	0.00	8.19	0.55	4.22	18.59	28.31	18.80	2.77	15.51	0.00	2.46	0.58	0.39
Cosuppresion 2.3	0.00	8.71	0.47	3.48	19.21	30.06	19.49	2.03	13.78	0.00	2.15	0.63	0.39
Cosuppresion 3.1	0.06	7.57	0.50	3.83	32.24	20.00	15.66	2.06	15.65	0.34	1.85	0.23	0.90
Cosuppresion 4.1	0.00	7.29	0.43	3.55	30.26	21.17	17.06	2.01	16.08	0.00	1.92	0.25	0.79
Cosuppresion 4.2	0.08	8.02	0.53	3.62	33.04	20.04	15.68	1.80	14.72	0.00	1.88	0.58	0.92
Cosuppresion 4.3	0.07	8.35	0.54	3.85	30.02	21.72	16.78	2.01	14.25	0.00	1.92	0.49	0.78
Cosuppresion 4.4	0.06	6.98	0.53	3.62	43.38	13.24	12.77	1.74	15.37	0.30	1.67	0.33	1.67
Cosuppresion 4.5	0.13	7.84	0.52	3.76	33.76	18.16	16.21	1.89	14.96	0.35	1.85	0.57	0.98
Cosuppresion 4.6	0.11	8.18	0.32	3.58	19.72	29.19	20.26	2.04	13.92	0.29	1.84	0.55	0.40
Cosuppresion 4.7	0.11	7.88	0.39	3.75	27.40	22.85	17.44	2.08	15.29	0.00	2.04	0.76	0.68
Cosuppresion 4.8	0.13	7.56	0.41	3.46	32.27	20.50	15.45	1.90	15.47	0.00	2.02	0.83	0.90
Cosuppresion 4.9	0.09	7.46	0.29	3.75	36.11	16.96	15.74	1.92	15.38	0.31	1.74	0.25	1.10
Cosuppresion 5.1	0.10	7.68	0.34	3.88	36.00	16.77	15.38	1.90	15.44	0.32	1.82	0.36	1.12

5

Analysis of the results indicates that transgenic plants harboring a CoP construct (indicated as "hairpin x.x" in the table) have a higher frequency of plants with oil wherein the increase in oleic acid and concomitant decrease in linolenic and linoleic acid is significant than in transgenic plants harboring cosuppression constructs.. Moreover the absolute levels of increase, respectively decrease are higher respectively lower than in transgenic plants harboring cosuppression constructs.

Example 7. Modifying oil profile using CoP constructs in *Brassica*.

The T-DNA vector harboring the chimeric gene encoding the CoP construct described in Example 6 is introduced in *Brassica* oilseed rape. Seeds harvested from the transgenic *Brassica sp.* are crashed and oil extracted and the composition of the fatty acids in the oil is analyzed.

Oil from transgenic *Brassica sp.* harboring the CoP construct have significantly increased oleic acid content and decreased linoleic and linolenic acid content. A T-DNA vector harboring a chimeric gene encoding a CoP construct similar to the one described in Example 6, but wherein the sequence of the sense and antisense region corresponding to the $\Delta 12$ desaturase encoding ORF is based on a homologous ORF from *Brassica spp.* is constructed and introduced in *Brassica* oilseed rape.

The sequence of *Brassica spp* ORFs homologous to $\Delta 12$ desaturase encoding ORF from *Arabidopsis* are available from Genbank database under Accession nrs AF042841 and AF124360.

Seeds harvested from the transgenic *Brassica sp.* are crashed and oil extracted and the composition of the fatty acids in the oil is analyzed. Oil from transgenic *Brassica sp.* harbouring the CoP construct have significantly increased oleic acid content and decreased linoleic and linolenic acid content.

Example 8. Suppression of an endogenous rust resistance gene in flax.

A CoP construct for suppression of the endogenous rust resistance gene was made consisting of

1. a CaMV35S promoter; operably linked to
2. part of an endogenous rust resistance gene (n) from flax (about 1500 bp long) in the sense orientation; ligated to
3. a similar part of the endogenous rust resistance gene from flax (about 1450 bp long) in antisense orientation so that a perfect inverted repeat without spacer sequence is generated wherein each repeat is about 1450 bp long; followed by
4. a nos terminator.

Plain antisense constructs were made using a similar fragment as described sub 3 above inserted between a CaMV35S promoter and a nos terminator.

Flax plants containing the n gene (which gives resistance to a strain of flax rust) were transformed by these CoP and antisense constructs. If suppression occurs, the plants become susceptible to the rust strain. If the construct has no effect, the transformed plants remain resistant to the rust strain.

Results

ngc-b sense/antisense	3 suppressed out of 7
ngc-b antisense	0 suppressed out of 12

While the invention has been described and illustrated herein by references to various specific material, procedures and examples, it is understood that the invention is not restricted to the particular material, combinations of material, and procedures selected for that purpose. Numerous variations of such details can be implied and will be appreciated by those skilled in the art.

REFERENCES

- An et al. (1996) *The Plant Cell* 8, 15-30
- Barry et al. (1993) *Proc Natl Acad Sci* 90, 4557-4561
- Baulcombe (1996) *Plant Cell* 8, 1833-1844
- 5 Braun and Hemenway (1992) *Plant Cell* 4, 735-744
- Brederode et al. (1995) *Virology* 207, 467-474
- Carr et al. (1992) *Mol. Plant-Microb. Interact.* 5, 397-404
- Christensen and Quail (1996) *Transgenic Research* 5, 213-218
- Croy Plant Molecular Biology Labfax (1993) BIOS Scientific Publications Ltd (UK)
- 10 and Blackwell Scientific Publications, UK.
- de Carvalho Niebel et al. (1995) *Plant Cell* 7, 347-358
- English et al. (1996) *Plant Cell* 8, 179-188
- Fire et al. (1998) *Nature* 391, 806-811
- Fromm et al. (1990) *Bio/Technology* 8: 833
- 15 Gleave, (1992) *Plant Mol. Biol.* 20: 1203-1207
- Goodwin et al. (1996) *Plant Cell* 8, 95-105
- Gordon-Kamm et al. (1990) *The Plant Cell* 2: 603
- Harpster et al. (1988) *Mol. Gen. Genet.* 212, 182-190
- Hobbs et al. (1990) *Plant Mol. Biol.* 15, 851-864
- 20 Hudspeth et al. (1989) *Plant Mol Biol* 12: 579-589
- Ingelbrecht et al. (1994) *Proc. Natl. Acad. Sci. USA* 91, 10502-10506
- Jefferson et al. (1987) *EMBO J.* 6, 3901-3907
- Kawchuck et al. (1991) *Molecular plant-microbe interactions* 4, 247-253.
- Keil et al. (1989) *EMBO J.* 8: 1323-1330
- 25 Keller et al. (1988) *EMBO J.* 7: 3625-3633
- Keller et al. (1989) *Genes Devel.* 3: 1639-1646
- Landsman et al. (1988) *Mol Gen Genet* 214, 68-73
- Lindbo & Dougherty (1992a) *Mol. Plant Micr. Int* 5, 144-153
- Lindbo & Dougherty (1992b) *Virology* 189, 725-733
- 30 Lindbo et al. (1993) *Plant Cell* 5, 1749-1759
- Longstaff et al. (1993) *EMBO J.* 12, 379-386

MacDonald et al. (1991) *Nucl. Acids Res.* 19, 5575-5581

Metzlaff et al. (1997) *Cell* 88, 845-854

Meyer et al. (1987) *Nature* 330: 677

Mueller et al. (1995) *Plant J.* 7, 1001-1003

5 Ohta et al. (1990) *Plant Cell. Physiol.* 31, 805-813

Pang et al. (1996) *Plant J.* 9, 899-909

Peleman et al. 1989 *Gene* 84: 359-369

Powell-Abel et al. (1986) *Science* 232, 738-743

Powell et al. (1990) *Virology* 175, 124-130

10 Que et al. (1998) *The Plant Journal* 13, 401-409

Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition,
Cold Spring Harbor Laboratory Press, NY

Sanford and Johnston, (1985) *J. Theor. Biol.* 113, 395-405

Schiebel et al. (1993a) *Journal of Biological Chemistry* 268: 11851-11857

15 Schiebel et al. (1993b) *Journal of Biological Chemistry* 268: 11858-11867

Smith et al. (1994) *Plant Cell* 6, 1441-1453

Stalberg et al., *Plant Molecular Biol.* 23, 671-683

Stam et al. (1997) *Ann. Botan.* 79, 3-12

Wagner and Sun (1998) *Nature* 391, 744-745

20 Wang et al. (1997) *Journal of Genetics and Breeding* 3

Wilbur and Lipmann (1983) *Proc. Nat. Acad. Sci. U.S.A.* 80: 726

Zheng et al. (1991) *Plant Physiol.* 97, 832-835

Zuker and Stiegler (1981) *Nucl. Acids Res.* 9, 133-148

WHAT IS CLAIMED IS:

1. A method for reducing the phenotypic expression of a nucleic acid of interest, which is normally capable of being expressed in a eucaryotic cell, comprising the step of introducing a chimeric DNA comprising the following operably linked parts:

- a) a promoter, operative in said eucaryotic cell;
- b) DNA region, which when transcribed, yields an RNA molecule comprising an RNA region capable of forming an artificial stem-loop structure, wherein one of the annealing RNA sequences of the stem-loop structure comprises a sequence, essentially similar to at least part of the nucleotide sequence of said nucleic acid of interest, and wherein the second of said annealing RNA sequences comprises a sequence essentially similar to at least part of the complement of at least part of said nucleotide sequence of said nucleic acid of interest; and optionally
- c) a DNA region involved in transcription termination and polyadenylation.

2. A method for reducing the phenotypic expression of a nucleic acid of interest, which is normally capable of being expressed in a eucaryotic cell, comprising the step of introducing a chimeric DNA comprising the following operably linked parts:

- a) a promoter, operative in said eucaryotic cell;
- b) a DNA region, which when transcribed, yields an RNA molecule with a nucleotide sequence comprising
 - i. a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of said nucleic acid of interest; and
 - ii. an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;

wherein the RNA is capable of forming an artificial hairpin RNA structure with a double stranded RNA stem by base-pairing between the regions with sense and antisense nucleotide sequence such that at least said 10 consecutive

nucleotides of the sense sequence basepair with said 10 consecutive nucleotides of the antisense sequence; and optionally

c) a DNA region involved in transcription termination and polyadenylation.

5 3. The method of claim 2, wherein said RNA molecule further comprises a spacer nucleotide sequence located between said sense and said antisense nucleotide sequence.

10 4. The method of claim 2, wherein said sense nucleotide sequence comprises at least about 550 consecutive nucleotides having between 75% and 100% sequence identity with at least part of the nucleotide sequence of said nucleic acid.

15 5. The method of claim 2, wherein said nucleic acid of interest is a gene integrated in the genome of said eucaryotic cell.

6. The method of claim 5, wherein said gene is an endogenous gene.

7. The method of claim 5, wherein said gene is a foreign transgene.

20 8. The method of claim 2, wherein said chimeric DNA is stably integrated in the genome of the DNA.

25 9. The method of claim 2, wherein said nucleic acid of interest is comprised in the genome of an infecting virus.

10. The method of claim 9, wherein said infecting virus is an RNA virus.

30 11. The method according to claim 1, wherein said eucaryotic cell is a plant cell.

12. The method of claim 11, wherein said plant cell is comprised within a plant.

13. A method for reducing the phenotypic expression of a nucleic acid of interest, which is normally capable of being expressed in a eucaryotic cell comprising the step of introducing a chimeric RNA molecule comprising at least one RNA region with a nucleotide sequence comprising

- i. a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and
- ii. an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;

wherein said RNA region is capable of forming an artificial hairpin RNA structure with a double stranded RNA stem by base-pairing between the regions with sense and antisense nucleotide sequence such that at least said 10 consecutive nucleotides of the sense sequence basepair with said 10 consecutive nucleotides of the antisense sequence.

14. A method for reducing the gene expression of a gene of interest in plant cells, said method comprising the step of introducing a first and second chimeric DNA, linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of the transgenic plant cells; wherein said first chimeric DNA comprises the following operably linked parts:

- a) a plant-expressible promoter;
- b) a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of said gene of interest; and optionally
- c) a DNA region involved in transcription termination and polyadenylation functioning in plant cells; and

wherein said second chimeric DNA comprises the following operably linked parts:

- a) a plant-expressible promoter;
- b) a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence; and optionally
- c) a DNA region involved in transcription termination and polyadenylation functioning in plant cells;

wherein said sense and antisense RNA molecules are capable of forming a double stranded RNA by base-pairing between the regions which are complementary.

15. A method for obtaining a virus resistant eucaryotic organism, said method comprising the steps of :

- 1) providing the cells of said organism with a first and second chimeric DNA wherein said first chimeric DNA comprises the following operably linked parts:
 - a) a promoter operative in said cells;
 - b) a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the genome of a virus, capable of infecting said organism; and optionally
 - c) a DNA region involved in transcription termination and polyadenylation functioning in said cells; and

wherein said second chimeric DNA comprises the following operably linked parts:

- a) a promoter operative in said cells;
- b) a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the

complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence; and optionally

- c) a DNA region involved in transcription termination and polyadenylation functioning in said cells; and

5 wherein said sense and antisense RNA are capable of forming a double stranded RNA by base-pairing between the regions which are complementary.

16. The method according to claim 15, wherein said organism is a plant

10 17. The method of claim 16, wherein said cells of said plants are provided with said first and second chimeric DNA by crossing parent plants comprising either said first or said second chimeric DNA.

15 18. The method of claim 16, wherein said cells of said plants are provided with said first and second chimeric DNA by transforming a plant cell with said first and second chimeric DNA, and regenerating a plant from said transformed plant cell.

19. The method of claim 16, wherein said first and second chimeric DNA are integrated separately in said nuclear genome of said plant cell.

20 20. The method of claim 16, wherein said first and second chimeric DNA are linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of the transgenic plant cells.

25 21. A method for identifying a phenotype associated with the expression of a nucleic acid of interest in a eucaryotic cell, said method comprising

- a. selecting within said nucleotide sequence of interest, a target sequence of at least 10 consecutive nucleotides;
 - b. designing a sense nucleotide sequence corresponding to the length of
- 30 the selected target sequence and which has a sequence identity of at least about 75% to about 100% with said selected target sequence;

- c. designing an antisense nucleotide sequence which:
 - i) has a sequence identity of at least about 75% to about 100% with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;
 - ii) comprises a stretch of at least about 10 consecutive nucleotides with 100% sequence identity to the complement of a part of said sense nucleotide sequence;
- d. introducing an RNA molecule comprising both said sense and antisense nucleotide sequences into a suitable eucaryotic host cell comprising the nucleic acid including the nucleotide sequence with hitherto unidentified phenotype; and
- e. observing the phenotype by a suitable method.

22. A eucaryotic cell, comprising a nucleic acid of interest, which is normally capable of being phenotypically expressed, further comprising a chimeric DNA molecule comprising the following operably linked parts:

- a) a promoter, operative in said eucaryotic cell;
- b) a DNA region, which when transcribed, yields an RNA molecule with at least one RNA region with a nucleotide sequence comprising
 - i. a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and
 - ii. an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;

wherein the RNA is capable of forming an artificial hairpin RNA structure with a double stranded RNA stem by base-pairing between the regions with sense and antisense nucleotide sequence; and optionally

c) a DNA region involved in transcription termination and polyadenylation. wherein the phenotypic expression of said nucleic acid of interest is significantly reduced.

23. A eucaryotic cell, comprising a nucleic acid of interest, which is normally capable of being phenotypically expressed, further comprising a chimeric RNA molecule comprising at least one RNA region with a nucleotide sequence comprising

- i. a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the nucleic acid of interest; and
- ii. an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;

wherein said RNA is capable of forming an artificial hairpin RNA with a double stranded RNA region by base-pairing between the regions with sense and antisense nucleotide sequence such that at least said 10 consecutive nucleotides of the sense sequence basepair with said 10 consecutive nucleotides of the antisense sequence.

24. A eucaryotic cell, comprising a gene of interest, which is normally capable of being phenotypically expressed, further comprising a first and second chimeric DNA, linked on one recombinant DNA such that both chimeric DNAs are integrated together in the nuclear genome of said eucaryotic cell; wherein said first chimeric DNA comprises the following operably linked parts:

- a) a promoter operative in said eucaryotic cell
- b) a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of said gene of interest; and optionally

c) a DNA region involved in transcription termination and polyadenylation; and wherein said second chimeric DNA comprises the following operably linked parts:

- a) a promoter operative in said eucaryotic cell;
- b) a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence; and optionally

c) a DNA region involved in transcription termination and polyadenylation; wherein said sense and antisense RNA molecules are capable of forming a double stranded RNA region by base-pairing between the regions which are complementary.

25. The eucaryotic cell of claim 22, which is a plant cell.

26. A plant comprising the plant cell of claim 25.

27. A virus resistant plant, comprising a first and second chimeric DNA integrated in the nuclear genome its cells, wherein said first chimeric DNA comprises the following operably linked parts:

- a) a plant-expressible promoter;
- b) a first DNA region capable of being transcribed into a sense RNA molecule with a nucleotide sequence comprising a sense nucleotide sequence of at least 10 consecutive nucleotides having between 75 and 100% sequence identity with at least part of the nucleotide sequence of the genome of a virus, capable of infecting said plant; and optionally
- c) a DNA region involved in transcription termination and polyadenylation functioning in plant cells; and

wherein said second chimeric DNA comprises the following operably linked parts:

- a) a plant-expressible promoter;

- b) a second DNA region capable of being transcribed into an antisense RNA molecule with a nucleotide sequence comprising an antisense nucleotide sequence including at least 10 consecutive nucleotides, having between about 75% to about 100% sequence identity with the complement of said at least 10 consecutive nucleotides of said sense nucleotide sequence;
- c) optionally, a DNA region involved in transcription termination and polyadenylation functioning in plant cells; and

wherein said sense and antisense RNA are capable of forming a double stranded RNA region by base-pairing between the regions which are complementary.

28. The plant of claim 27, wherein said first and second chimeric DNA are integrated in one locus in the nuclear genome.

29. The plant of claim 27, wherein said first and second chimeric DNA are integrated in different loci in the nuclear genome.

30. A method for modifying the fatty acid profile in oil from a plant, said method comprising the step of introducing a chimeric DNA into the cells of said plant, said chimeric DNA comprising the following operably linked parts:

- a). a plant-expressible promoter
- b). a DNA region, which when transcribed yields an RNA molecule comprising an RNA region capable of forming an artificial stem-loop structure, wherein one of the annealing RNA sequences of the stem-loop structure comprises a nucleotide sequence essentially similar to at least part of the nucleotide sequence of a $\Delta 12$ desaturase encoding open reading frame, and wherein the second of said annealing RNA sequences comprises a sequence essentially similar to at least part of the complement of at least part of the nucleotide sequence of said $\Delta 12$ desaturase encoding open reading frame; and optionally;
- c) a DNA region involved in transcription termination and polyadenylation.

31. The method of claim 30, wherein said DNA region comprises the nucleotide sequence of SEQ ID No 6.

32. The method of claim 30, wherein said modifying of the fatty acid profile, comprises increasing the oleic acid content.

33. The method of claim 30, wherein said plant expressible promoter is a seed-specific promoter.

34. The method of claim 30, wherein said plant is oilseed rape.

35. A plant producing oil with modified fatty acid profile, said plant comprising a chimeric DNA, said chimeric DNA comprising the following operably linked parts:

- a). a plant-expressible promoter
- b). a DNA region, which when transcribed yields an RNA molecule comprising an RNA region capable of forming an artificial stem-loop structure, wherein one of the annealing RNA sequences of the stem-loop structure comprises a nucleotide sequence essentially similar to at least part of the nucleotide sequence of a $\Delta 12$ desaturase encoding open reading frame, and wherein the second of said annealing RNA sequences comprises a sequence essentially similar to at least part of the complement of at least part of the nucleotide sequence of said $\Delta 12$ desaturase encoding open reading frame; and optionally;
- c). a DNA region involved in transcription termination and polyadenylation.

36. The plant of claim 35, wherein said DNA region comprises the nucleotide sequence of SEQ ID No 6.

37. The plant of claim 35, wherein said plant expressible promoter is a seed-specific promoter.

38. The plant of claim 35, wherein said plant is oilseed rape.

ABSTRACT OF THE DISCLOSURE

Methods and means are provided for reducing the phenotypic expression of a nucleic acid of interest in eucaryotic cells, particularly in plant cells, by introducing chimeric genes encoding sense and antisense RNA molecules directed towards the target nucleic acid, which are capable of forming a double stranded RNA region by base-pairing between the regions with sense and antisense nucleotide sequence or by introducing the RNA molecules themselves. Preferably, the RNA molecules comprises simultaneously both sense and antisense nucleotide sequence.

5




Figure 1A: Schematic representation of the chimeric genes used in Example 1





Figure 1B: Schematic representation of the chimeric genes used in Examples 2 and 4

Figure 2



**COMBINED DECLARATION AND POWER OF ATTORNEY
FOR UTILITY PATENT APPLICATION**

Attorney's Docket No.
021565-040

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I BELIEVE I AM THE ORIGINAL, FIRST AND SOLE INVENTOR (if only one name is listed below) OR AN ORIGINAL, FIRST AND JOINT INVENTOR (if more than one name is listed below) OF THE SUBJECT MATTER WHICH IS CLAIMED AND FOR WHICH A PATENT IS SOUGHT ON THE INVENTION ENTITLED:

METHODS AND MEANS FOR OBTAINING MODIFIED PHENOTYPES

the specification of which

(check one)

☐ is attached hereto;

☒ was filed on April 8, 1998 as

Application No. 09/056,767

and was amended on _____;
(if applicable)

I HAVE REVIEWED AND UNDERSTAND THE CONTENTS OF THE ABOVE-IDENTIFIED SPECIFICATION, INCLUDING THE CLAIMS, AS AMENDED BY ANY AMENDMENT REFERRED TO ABOVE;

I ACKNOWLEDGE THE DUTY TO DISCLOSE TO THE OFFICE ALL INFORMATION KNOWN TO ME TO BE MATERIAL TO PATENTABILITY AS DEFINED IN TITLE 37, CODE OF FEDERAL REGULATIONS, Sec. 1.56 (as amended effective March 16, 1992);

I do not know and do not believe the said invention was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to said application; that said invention was not in public use or on sale in the United States of America more than one year prior to said application; that said invention has not been patented or made the subject of an inventor's certificate issued before the date of said application in any country foreign to the United States of America on any application filed by me or my legal representatives or assigns more than twelve months prior to said application;

I hereby claim foreign priority benefits under Title 35, United States Code Sec. 119 and/or Sec. 365 of any foreign application(s) for patent or inventor's certificate as indicated below and have also identified below any foreign application for patent or inventor's certificate on this invention having a filing date before that of the application(s) on which priority is claimed:

COPY

DOCKETED
filed 10/23/98

COMBINED DECLARATION AND POWER OF ATTORNEY			Attorney's Docket No. 021565-040																																																																															
COUNTRY/INTERNATIONAL	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED																																																																															
			YES _ NO _																																																																															
			YES _ NO _																																																																															
<p>I hereby appoint the following attorneys and agent(s) to prosecute said application and to transact all business in the Patent and Trademark Office connected therewith and to file, prosecute and to transact all business in connection with international applications directed to said invention:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;">William L. Mathis</td> <td style="width: 10%;">17,337</td> <td style="width: 33%;">Robert G. Mukai</td> <td style="width: 10%;">28,531</td> <td style="width: 33%;">Bruce J. Boggs, Jr.</td> <td style="width: 10%;">32,344</td> </tr> <tr> <td>Peter H. Smolka</td> <td>15,913</td> <td>George A. Hovanec, Jr.</td> <td>28,223</td> <td>William H. Benz</td> <td>25,952</td> </tr> <tr> <td>Robert S. Swecker</td> <td>19,885</td> <td>James A. LaBarre</td> <td>28,632</td> <td>Peter K. Skiff</td> <td>31,917</td> </tr> <tr> <td>Platon N. Mandros</td> <td>22,124</td> <td>E. Joseph Gess</td> <td>28,510</td> <td>Richard J. McGrath</td> <td>29,195</td> </tr> <tr> <td>Benton S. Duffett, Jr.</td> <td>22,030</td> <td>R. Danny Huntington</td> <td>27,903</td> <td>Manthaw L. Schneider</td> <td>32,814</td> </tr> <tr> <td>Joseph R. Magnone</td> <td>24,239</td> <td>Eric H. Weisblatt</td> <td>30,505</td> <td>Michael G. Savage</td> <td>32,596</td> </tr> <tr> <td>Norman H. Stepno</td> <td>22,716</td> <td>James W. Peterson</td> <td>26,057</td> <td>Gerald F. Swiss</td> <td>30,113</td> </tr> <tr> <td>Ronald L. Grudziecki</td> <td>24,970</td> <td>Teresa Stanek Rea</td> <td>30,427</td> <td>Michael J. Ure</td> <td>33,089</td> </tr> <tr> <td>Frederick G. Michaud, Jr.</td> <td>26,003</td> <td>Robert E. Krebs</td> <td>25,885</td> <td>Charles F. Wieland III</td> <td>33,096</td> </tr> <tr> <td>Alan E. Kopecki</td> <td>25,813</td> <td>Robert M. Schulman</td> <td>31,196</td> <td>Bruce T. Wieder</td> <td>33,815</td> </tr> <tr> <td>Regis E. Sluter</td> <td>26,999</td> <td>William C. Rowland</td> <td>30,888</td> <td>Todd R. Walters</td> <td>34,040</td> </tr> <tr> <td>Samuel C. Miller, III</td> <td>27,360</td> <td>T. Gene Dillahunty</td> <td>25,423</td> <td></td> <td></td> </tr> <tr> <td>Ralph L. Freeland, Jr.</td> <td>16,110</td> <td>Patrick C. Keane</td> <td>32,858</td> <td></td> <td></td> </tr> </table>					William L. Mathis	17,337	Robert G. Mukai	28,531	Bruce J. Boggs, Jr.	32,344	Peter H. Smolka	15,913	George A. Hovanec, Jr.	28,223	William H. Benz	25,952	Robert S. Swecker	19,885	James A. LaBarre	28,632	Peter K. Skiff	31,917	Platon N. Mandros	22,124	E. Joseph Gess	28,510	Richard J. McGrath	29,195	Benton S. Duffett, Jr.	22,030	R. Danny Huntington	27,903	Manthaw L. Schneider	32,814	Joseph R. Magnone	24,239	Eric H. Weisblatt	30,505	Michael G. Savage	32,596	Norman H. Stepno	22,716	James W. Peterson	26,057	Gerald F. Swiss	30,113	Ronald L. Grudziecki	24,970	Teresa Stanek Rea	30,427	Michael J. Ure	33,089	Frederick G. Michaud, Jr.	26,003	Robert E. Krebs	25,885	Charles F. Wieland III	33,096	Alan E. Kopecki	25,813	Robert M. Schulman	31,196	Bruce T. Wieder	33,815	Regis E. Sluter	26,999	William C. Rowland	30,888	Todd R. Walters	34,040	Samuel C. Miller, III	27,360	T. Gene Dillahunty	25,423			Ralph L. Freeland, Jr.	16,110	Patrick C. Keane	32,858		
William L. Mathis	17,337	Robert G. Mukai	28,531	Bruce J. Boggs, Jr.	32,344																																																																													
Peter H. Smolka	15,913	George A. Hovanec, Jr.	28,223	William H. Benz	25,952																																																																													
Robert S. Swecker	19,885	James A. LaBarre	28,632	Peter K. Skiff	31,917																																																																													
Platon N. Mandros	22,124	E. Joseph Gess	28,510	Richard J. McGrath	29,195																																																																													
Benton S. Duffett, Jr.	22,030	R. Danny Huntington	27,903	Manthaw L. Schneider	32,814																																																																													
Joseph R. Magnone	24,239	Eric H. Weisblatt	30,505	Michael G. Savage	32,596																																																																													
Norman H. Stepno	22,716	James W. Peterson	26,057	Gerald F. Swiss	30,113																																																																													
Ronald L. Grudziecki	24,970	Teresa Stanek Rea	30,427	Michael J. Ure	33,089																																																																													
Frederick G. Michaud, Jr.	26,003	Robert E. Krebs	25,885	Charles F. Wieland III	33,096																																																																													
Alan E. Kopecki	25,813	Robert M. Schulman	31,196	Bruce T. Wieder	33,815																																																																													
Regis E. Sluter	26,999	William C. Rowland	30,888	Todd R. Walters	34,040																																																																													
Samuel C. Miller, III	27,360	T. Gene Dillahunty	25,423																																																																															
Ralph L. Freeland, Jr.	16,110	Patrick C. Keane	32,858																																																																															
<p>and: _____</p>																																																																																		
<p>Address all correspondence to: <u>R. Danny Huntington</u> <u>BURNS, DOANE, SWECKER & MATHIS, L.L.P.</u> <u>P.O. Box 1404</u> <u>Alexandria, Virginia 22313-1404</u></p>																																																																																		
<p>Address all telephone calls to: <u>R. Danny Huntington</u> at (703) 836-6620.</p>																																																																																		
<p>I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.</p>																																																																																		
FULL NAME OF SOLE OR FIRST INVENTOR		SIGNATURE		DATE																																																																														
Peter Michael WATERHOUSE		<i>Peter Waterhouse</i>		12/6/98																																																																														
RESIDENCE			CITIZENSHIP																																																																															
Canberra ACT 2602, Australia			Australian and United Kingdom																																																																															
POST OFFICE ADDRESS																																																																																		
5 Banjine Street, O. Connor, Canberra, ACT 2602 Australia																																																																																		
FULL NAME OF SECOND JOINT INVENTOR, IF ANY		SIGNATURE		DATE																																																																														
Ming-Bo WANG		<i>Mingbo Wang</i>		12/6/98																																																																														
RESIDENCE			CITIZENSHIP																																																																															
Canberra ACT 2612, Australia			Peoples Republic of China																																																																															
POST OFFICE ADDRESS																																																																																		
3/14 Masson Street, Turner, Canberra, ACT 2612 Australia																																																																																		
FULL NAME OF THIRD JOINT INVENTOR, IF ANY		SIGNATURE		DATE																																																																														
Michael Wayne Graham		<i>MW Graham</i>		16/6/98																																																																														
RESIDENCE			CITIZENSHIP																																																																															
St. Lucia QLD 4067, Australia			Australia																																																																															
POST OFFICE ADDRESS																																																																																		
42 Raglan Street, St. Lucia QLD 4067, Australia																																																																																		

COPY

SEQUENCE LISTING

<110> Waterhouse, Peter M

Wang, Ming-Bo

Graham, Michael W

Commonwealth Scientific and Industrial Research Or

<120> Methods and means for obtaining modified phenotypes

<130> echidna

<140>

<141>

<150> US SN 09/056,767

<151> 1998-03-08

<150> US SN 09/127735

<151> 1998-08-03

<160> 7

<170> PatentIn Ver. 2.0

<210> 1

<211> 854

<212> DNA

<213> Potato virus Y

<220>

<223> fragment of the NIa ORF

<400> 1

```

aagctttgaa gattgatttg atgccacata acccactcaa aatttgtgac aaaacaaatg 60
gcattgccaa atttcctgag agagagttcg agctaaggca gactgggcca gctgtagaag 120
tcgacgtgaa ggacatacca gcacaggagg tggaacatga agctaaatcg ctcatgagag 180
gcttgagaga cttcaacca attgccaaa cagtttgtag gctgaaagta tctgttgaat 240
atgggacatc agagatgtac ggttttggat ttggagcgta cataatagcg aaccaccatt 300
tggtcaggag ttataatggt tccatggagg tacgatccat gcacggtaca ttcagggtaa 360
agaatctaca cagtttgagc gttctgccaa ttaaaggtag ggacatcatc ctcattaaaa 420
tgccaaaaga tttccctgtc tttccacaga aattgcattt ccgagctcct acacagaacg 480
aaagaatttg tttagttgga accaactttc aggagaagta tgcacgtcgc atcatcacag 540
aagcaagcac tacttacaat ataccaggca gcacattctg gaagcattgg attgaaacag 600
ataatggaca ctgtggacta ccagtggtga gcaactgccg tggatgtcta gtcggaattc 660
acagtttggc aaacaatgca cacaccacga actactactc agccttcgat gaagattttg 720
aaagcaagta cctccgaacc aatgagcaca atgaatgggt caagtcttgg atttataatc 780
cagacacagt gttgtggggc ccgttgaaac ttaaagacag cactcctaaa gggttattta 840
aaacaacaaa gctt
854

```

<210> 2
 <211> 2186
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: coding region
 of the Gusd CoP construct

<220>
 <221> misc_structure
 <222> ()..
 <223> deficient Gus coding region

<220>
 <221> misc_feature
 <222> ()..(2186)
 <223> antisense to the 5' end of the Gus coding region

<400> 2
 atggtacgtc ctgtagaaac cccaacccgt gaaatcaaaa aactcgacgg cctgtgggca 60
 ttcagtctgg atcgcgaaaa ctgtggaatt gatcagcgtt ggtgggaaag cgcgttacia 120
 gaaagccggg caattgctgt gccaggcagt tttaacgac agttcgccga tgcagatatt 180
 cgtaattatg cgggcaacgt ctggtatcag cgcgaaagtct ttataccgaa aggttgggca 240
 ggccagcgta tctgtctgctg tttcgatgctg gtcactcatt acggcaaagt gtgggtcaat 300
 aatcaggaag tgatggagca tcagggcggc tatacgccat ttgaagccga tgcacgccc 360
 tatgttattg cggggaaaag tgtacgtatc accgtttgtg tgaacaacga actgaactgg 420
 cagactatcc cgcggggaat ggtgattacc gacgaaaacg gcaagaaaaa gcagtcttac 480
 ttccatgatt tctttaacta tgccggaatc catcgacgct taatgctcta caccagccg 540
 aacacctggg tggacgatat ctaccgcgtt cgcgctcgga tccggtcagt ggcagtgaag 600
 ggcgaaacgt tcttgattaa ccacaaaccg ttctacttta ctggcttttg tctcatgaa 660
 gatgctggact tgcgtggcaa aggattcgat aacgtgctga tgggtgcacg ccacgcatta 720
 atggactgga ttggggccaa ctctaccgt accctgcatt acccttacgc tgaagagatg 780
 ctcgactggg cagatgaaca tggcatcgtg gtgattgatg aaactgctgc tgcggtctt 840
 aacctctctt taggcattgg tttcgaagcg ggcaacaagc cgaaagaact gtacagcgaa 900
 gaggcagtca acggggaaac tcagcaagcg cacttacagg cgattaaaga gctgatagcg 960
 cgtgacaaaa accacccaag cgtgggtgat tggagtattg ccaacgaacc ggatacccg 1020
 ccgcaagggtg cacgggaata tttcgcgcca ctggcggaag caacgcgtaa actcgacccg 1080
 acgcgtccga tcacctgctg caatgtaatg ttctgcgacg ctacacccga taccatcagc 1140
 gatctctttg atgtgctgtg cctgaaccgt tattacggat ggtatgtcca aagcggcgat 1200
 ttggaaacgg cagagaaggt actggaaaaa gaacttcttg cctggcagga gaaactgcat 1260
 cagccgatta tcatcaccga atacggcgtg gatacgtagg ccgggctgca ctcaatgtac 1320
 accgacatgt ggagtgaaga gtatcagtgt gcatggctgg atatgtatca ccgcgtcttt 1380
 gatcgctgca gcgcccgtcgt cgggtgaacag gtatggaatt tcgccgattt tgcgacctcg 1440
 caaggcatat tgcgcgttgg cggtaacaag aaaggatct tcaactcgca ccgcaaaccg 1500
 aagtcggcgg cttttctgct gcaaaaacgc tggactggca tgaacttcgg tgaaaaaccg 1560
 cagcagggag gcaaaacaatg aaacagacgc gtggttacag tcttgcgca catgcgtcac 1620

cacggtgata tcgtccaccc aggtgttcgg cgtgggtgtag agcatcacgt gcgatggatt 1680
 ccggcatagt taaagaaatc atggaagtaa gactgctttt tcttgccggt ttcgtcggta 1740
 atcaccattc ccggcgggat agtctgccag ttcagtctgt tgttcacaca aacggtgata 1800
 cgtacacttt tcccggcaat aacatacggc gtgacatcgg cttcaaattg cgtatagccg 1860
 ccctgatgct ccatcacttc ctgattattg acccacactt tgccgtaatg agtgaccgca 1920
 tcgaaacgca gcacgatacg ctggcctgcc caacctttcg gtataaagac ttcgcgctga 1980
 taccagacgt tgcccgcata attacgaata tctgcatcgg cgaactgacg gttaaaactg 2040
 cctggcacag caattgcccg gctttcttgt aacgcgcttt ccaccaacg ctgatcaatt 2100
 ccacagtttt cgcgatccag actgaatgcc cacaggccgt cgagtttttt gatttcacgg 2160
 gttggggttt ctacaggacg taccat 2186

<210> 3

<211> 208

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:5'UTR of
Johnson mosaic virus

<400> 3

cgccccgggc ccaacacaac acaacagaac ctacgtcaat tgattttatc aatcgcaaag 60
 ccttacaaag atcttcgcag tcgttcacat acagattcac cgaaccattc ttgttagctc 120
 tcgcacagag ataagcagga aaccatggca ggtgagtggg acacagtttg atagtaagag 180
 aaaccagagg aagactgcag gtacccgc 208

<210> 4

<211> 1150

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Subteranean
clover virus S4 promoter with S7 enhancer

<400> 4

aatctgcagc ggccgcttaa tagtaattat gattaattat gagataagag ttgttattat 60
 gcttatgagg aataaagaat gattaatatt gtttaatttt attccgcgaa gcggtgtgtt 120
 atgtttttgt tggagacatc acgtgactct cacgtgatgt ctccgcgaca ggctggcacg 180
 gggcttagta ttaccccggt ccggatcaga gacatttgac taaatattga cttggaataa 240
 tagcccttgg attagatgac acgtggacgc tcaggatctg tgatgctagt gaagcgctta 300
 agctgaacga atctgacgga agagcggaca tacgcacatg gattatggcc cacatgtcta 360
 aagtgtatct ctttacagct atattgatgt gacgtaagat gctttacttc gcttcgaagt 420
 aaagtaggaa attgctcgtc aagttattct tttctgaaag aaattattta attctaatta 480
 aattaaatga gtcgctataa atagtgtcga tgctgcctca catcgtattc ttcttcgcat 540
 cgtctgttct ggttttaagc gggatccagg cctcgagata tcggtacctt gttattatca 600
 ataaaagaat ttttattgtt attgtgttat ttggtaattt atgcttataa gtaattctat 660
 gattaattgt gaattattaa gactaatgag gataataatt gaatttgatt aaattaactc 720

tgcgaagcta tatgtctttc acgtgagagt cacgtgatgt ctccgcgaca ggctggcacg 780
 gggcttagta ttaccccggtg ccgggatcag agacatttga ctaaagtgtg acttgggaata 840
 atagcccttg gattagatga cacgtggacg ctcaggatct gtgatgctag tgaagcgctt 900
 aagctgaacg aatctgacgg aagagcggac aaacgcacat ggactatggc ccaactgcttt 960
 attaaagaag tgaatgacag ctgtctttgc ttcaagacga agtaaagaat agtggaaaac 1020
 gcgtaaagaa taagcgctact cagtacgctt cgtggcttta tataaatagt gcttcgtctt 1080
 attcttcggt gtatcatcaa cgaagaagtt aagctttgtt ctgcgtttta atgatcgatg 1140
 gccagtcgac 1150

<210> 5

<211> 1052

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: subterranean
 clover virus promoter S4 with S4 enhancer

<400> 5

ggatccaggc ctcgagatat cggtagcttg ttattatcaa taaaagaatt tttattgtta 60
 ttgtgttatt tggtaattha tgcttataag taattctatg attaattgtg aattattaag 120
 actaatgagg ataataattg aatttgatta aattaactct gcgaagctat atgtctttca 180
 cgtgagagtc acgtgatgtc tccgcgacag gctggcacgg ggcttagtat taccctgtgc 240
 cgggatcaga gacatttgac taaatgttga cttggaataa tagcccttgg attagatgac 300
 acgtggacgc tcaggatctg tgatgctagt gaagcgctta agctgaacga atctgacgga 360
 agagcggaca aacgcacatg gactatggcc cactgcttta ttaaagaagt gaatgacagc 420
 tgtctttgct tcaagacgaa gtaaagaata gtggaaaacg cgtggatcca ggcctcgaga 480
 tatcggtagc ttgttattat caataaaaaga atttttattg ttattgtgtt atttggtaat 540
 ttatgcttat aagtaattct atgattaatt gtgaattatt aagactaatg aggataataa 600
 ttgaatttga ttaaattaac tctgcgaagc tatatgtctt tcacgtgaga gtcacgtgat 660
 gtctccgcga caggctggca cggggcttag tattaccctg tgccgggacg agagacattt 720
 gactaaatgt tgacttgga taatagccct tggattagat gacacgtgga cgctcaggat 780
 ctgtgatgct agtgaagcgc ttaagctgaa cgaatctgac ggaagagcgg acaaacgcac 840
 atggactatg gccactgct ttattaaaga agtgaatgac agctgtcttt gcttcaagac 900
 gaagtaaaga atagtggaaa acgcgtaaag aataagcgta ctacgtacgc ttcgtggctt 960
 tatataaata gtgcttcgct ttattcttcg ttgtatcatc aacgaagaag ttaagctttg 1020
 ttctgcgttt taatgatcga tggccagtcg ac 1052

<210> 6

<211> 1583

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: coding
 sequence of the desaturase CoP construct

<220>

<221> misc_feature
 <222> (1)..(480)
 <223> corresponding to the 5' end of the
 delta12-desaturase (fad2) coding region, in sense
 orientation

<220>
 <221> misc_feature
 <222> (1101)..(1583)
 <223> corresponding to the 5' end of the
 delta12-desaturase (fad2) coding region, in anti
 sense orientation

<400> 6
 atcattatag cctcatgctt ctactacgtc gccaccaatt acttctctct cctccctcag 60
 cctctctctt acttggcttg gccactctat tgggcctgtc aaggctgtgt cctaactggg 120
 atctgggtca tagccacga atgcggtcac cacgcattca gcgactacca atggctggat 180
 gacacagttg gtcttatctt ccattccttc ctctctgtcc ctacttctc ctggaagtat 240
 agtcacgcc gtcaccattc caacactgga tccctcgaaa gagatgaagt atttgtccca 300
 aagcagaaat cagcaatcaa gtggtacggg aaatacctca acaaccctct tggacgcac 360
 atgatgttaa ccgctccagtt tgctctcggg tggcccttgt acttagcctt taacgtctct 420
 ggcagaccgt atgacgggtt cgcttgccat ttcttcccca acgctcccat ctacaatgac 480
 cgagaacgcc tccagatata cctctctgat gcgggtattc tagccgtctg ttttgggtct 540
 taccgttacg ctgctgcaca agggatggcc tcgatgatct gcctctacgg agtaccgctt 600
 ctgatagtga atgcgttcct cgtcttgatc acttacttgc agcactca tccctcgttg 660
 cctcactacg attcatcaga gtgggactgg ctccagggag ctttggctac cgtagacaga 720
 gactacggaa tcttgaacaa ggtgttccac aacattacag acacacacgt ggctcatcac 780
 ctgttctcga caatgccgca ttataacgca atggaagcta caaaggcgat aaagccaatt 840
 ctgggagact attaccagtt cgatggaaca ccgtggtatg tagcgatgta tagggaggca 900
 aaggagtgtg tctatgtaga accggacagg gaagggtgaca agaaagggtg gtactgggtac 960
 aacaataagt tatgagcatg atggtgaaga aattgtcgac ctttctcttg tctgtttgtc 1020
 ttttgttaaa gaagctatgc ttcgttttaa taatcttatt gtccattttg ttgtgttatg 1080
 acattttggc tgctcattat gttcagtaac atctaccctc gcaaccctt ctttaccggt 1140
 cgcttgggca gtatgccaga cggctctctg aatttccgat tcatgttccc ggtgggctcc 1200
 tgtttgacct gccaatgtga gtactacgca ggttctccca acaactccat aaagggcacg 1260
 gtgaactaac gactaaagac gaaaccctgt ttatgaagta gagaaagctc cctaggtcac 1320
 aaccttacca ctgccgctac tgatatgaag gtcctcttca ttccctgtc ctccttctt 1380
 accttctatt ctggttgaca cagtaggtcg gtaaccatca gcgacttacg caccactggc 1440
 gtaagcaccg gatactgggt ctatgggtcaa tcctgtgtcg gaactgtccg gggttatctc 1500
 ccggttcggt tcattctctc tccgactccc tcctctctct tcattaacca ccgctgcac 1560
 atcttcgtac tccgatatta cta 1583

<210> 7
 <211> 786
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: intron 2 of
the *Flaveria trinervia* purvate orthophosphate
dikinase

<400> 7

```

aagcttggtta aggaaataat tattttcttt tttcctttta gtataaaaata gttaagtgat 60
gttaattagt atgattataa taatatagtt gttataattg tgaaaaaata atttataaat 120
atattgttta cataaacaac atagtaatgt aaaaaaatat gacaagtgat gtgtaagacg 180
aagaagataa aagttgagag taagtatatt atttttaatg aatttgatcg aacatgtaag 240
atgatatact agcattaata tttgttttaa tcataatagt aattctagct ggtttgatga 300
attaaatatac aatgataaaa tactatagta aaaataagaa taaataaatt aaaataatat 360
ttttttatga ttaatagttt attatataat taaatatcta taccattact aaatatttta 420
gtttaaaagt taataaatat tttgttagaa attccaatct gcttgtaatt tatcaataaa 480
caaaatatta aataacaagc taaagtaaca aataatatca aactaataga aacagtaatc 540
taatgtaaca aaacataatc taatgctaata ataacaaagc gcaagatcta tcattttata 600
tagtattatt ttcaatcaac attcttatta atttctaaat aatacttgta gttttattaa 660
cttctaaatg gattgactat taattaaatg aattagtcga acatgaataa acaaggtaac 720
atgatagatc atgtcattgt gttatcattg atcttacatt tggattgatt acagttggga 780
aagctt                                     786

```

**COMBINED DECLARATION AND POWER OF ATTORNEY
FOR UTILITY PATENT APPLICATION**

Attorney's Docket No.
021565-040

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I BELIEVE I AM THE ORIGINAL, FIRST AND SOLE INVENTOR (if only one name is listed below) OR AN ORIGINAL, FIRST AND JOINT INVENTOR (if more than one name is listed below) OF THE SUBJECT MATTER WHICH IS CLAIMED AND FOR WHICH A PATENT IS SOUGHT ON THE INVENTION ENTITLED:

METHODS AND MEANS FOR OBTAINING MODIFIED PHENOTYPES

the specification of which

(check one)

☐ is attached hereto;

☒ was filed on April 8, 1998 as

Application No. 09/056,767

and was amended on _____;
(if applicable)

I HAVE REVIEWED AND UNDERSTAND THE CONTENTS OF THE ABOVE-IDENTIFIED SPECIFICATION, INCLUDING THE CLAIMS, AS AMENDED BY ANY AMENDMENT REFERRED TO ABOVE;

I ACKNOWLEDGE THE DUTY TO DISCLOSE TO THE OFFICE ALL INFORMATION KNOWN TO ME TO BE MATERIAL TO PATENTABILITY AS DEFINED IN TITLE 37, CODE OF FEDERAL REGULATIONS, Sec. 1.56 (as amended effective March 16, 1992);

I do not know and do not believe the said invention was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to said application; that said invention was not in public use or on sale in the United States of America more than one year prior to said application; that said invention has not been patented or made the subject of an inventor's certificate issued before the date of said application in any country foreign to the United States of America on any application filed by me or my legal representatives or assigns more than twelve months prior to said application;

I hereby claim foreign priority benefits under Title 35, United States Code Sec. 119 and/or Sec. 365 of any foreign application(s) for patent or inventor's certificate as indicated below and have also identified below any foreign application for patent or inventor's certificate on this invention having a filing date before that of the application(s) on which priority is claimed:

COPY

COMBINED DECLARATION AND POWER OF ATTORNEY

Attorney's Docket No.
021565-040

COUNTRY/INTERNATIONAL

APPLICATION NUMBER

DATE OF FILING
(day, month, year)

PRIORITY
CLAIMED

YES_ NO_

YES_ NO_

I hereby appoint the following attorneys and agent(s) to prosecute said application and to transact all business in the Patent and Trademark Office connected therewith and to file, prosecute and to transact all business in connection with international applications directed to said invention:

William L. Mathis 17,337
Peter H. Smolka 15,913
Robert S. Swecker 19,885
Platon N. Mandros 22,124
Benton S. Duffett, Jr. 22,030
Joseph R. Magnone 24,239
Norman H. Stepno 22,716
Ronald L. Grudziecki 24,970
Frederick G. Michaud, Jr. 26,003
Alan E. Kopecki 25,813
Regis E. Slutter 26,999
Samuel C. Miller, III 27,360
Ralph L. Freeland, Jr. 16,110

Robert G. Mukai 28,531
George A. Hovanec, Jr. 28,223
James A. LaBarre 28,632
E. Joseph Gess 28,510
R. Danny Huntington 27,903
Eric H. Weisblatt 30,505
James W. Peterson 26,057
Teresa Stanek Rea 30,427
Robert E. Krebs 25,885
Robert M. Schulman 31,196
William C. Rowland 30,888
T. Gene Dillahunty 25,423
Patrick C. Keane 32,858

Bruce J. Boggs, Jr. 32,344
William H. Benz 25,952
Peter K. Skiff 31,917
Richard J. McGrath 29,195
Matthew L. Schneider 32,814
Michael G. Savage 32,596
Gerald F. Swiss 30,113
Michael J. Ure 33,089
Charles F. Wieland III 33,096
Bruce T. Wieder 33,815
Todd R. Walters 34,040

and:

Address all correspondence to:

R. Danny Huntington
BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404

Address all telephone calls to: R. Danny Huntington at (703) 836-6620.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF SOLE OR FIRST INVENTOR

Peter Michael WATERHOUSE

SIGNATURE

Peter Waterhouse

DATE

12/6/98

RESIDENCE

Canberra ACT 2602, Australia

CITIZENSHIP

Australian and United Kingdom

POST OFFICE ADDRESS

5 Banjine Street, O. Connor, Canberra, ACT 2602 Australia

FULL NAME OF SECOND JOINT INVENTOR, IF ANY

Ming-Bo WANG

SIGNATURE

Mingbo Wang 2

DATE

12/6/98

RESIDENCE

Canberra ACT 2612, Australia

CITIZENSHIP

Peoples Republic of China

POST OFFICE ADDRESS

3/14 Masson Street, Turner, Canberra, ACT 2612 Australia

FULL NAME OF THIRD JOINT INVENTOR, IF ANY

Michael Wayne Graham

SIGNATURE

Michael Wayne Graham

DATE

16/6/98

RESIDENCE

St. Lucia QLD 4067, Australia

CITIZENSHIP

Australia

POST OFFICE ADDRESS

42 Raglan Street, St. Lucia QLD 4067, Australia

COPY